

High-accuracy comparison of numerical relativity simulations with post-Newtonian expansions

Michael Boyle,¹ Duncan A. Brown,^{1,2,3} Lawrence E. Kidder,⁴ Abdul H. Mroué,⁴
Harald P. Pfeiffer,¹ Mark A. Scheel,¹ Gregory B. Cook,⁵ and Saul A. Teukolsky⁴

¹*Theoretical Astrophysics 103-33, California Institute of Technology, Pasadena, CA 91125*

²*LIGO Laboratory, California Institute of Technology, Pasadena, California 91125*

³*Department of Physics, Syracuse University, Syracuse, New York, 13244*

⁴*Center for Radiophysics and Space Research, Cornell University, Ithaca, New York, 14853*

⁵*Department of Physics, Wake Forest University, Winston-Salem, North Carolina 27106*

(Dated: September 13, 2007)

Numerical simulations of 15 orbits of an equal-mass binary black hole system are presented. Gravitational waveforms from these simulations, covering more than 30 cycles and ending about 1.5 cycles before merger, are compared with those from quasi-circular zero-spin post-Newtonian (PN) formulae. The cumulative phase uncertainty of these comparisons is about 0.05 radians, dominated by effects arising from the small residual spins of the black holes and the small residual orbital eccentricity in the simulations. Matching numerical results to *generic* post-Newtonian time-domain Taylor approximants about 22 cycles before merger yields good agreement (within 0.05 radians) until 15 cycles before merger, but phase differences of several radians build up toward merger. Similarly, when matching to generic post-Newtonian approximants at a time just before merger, phase differences of several radians occur at early times. However, one specific post-Newtonian approximant, TaylorT4 at 3.5PN order, agrees much better with the numerical simulations, with accumulated phase differences of less than 0.05 radians over the 30-cycle waveform. Gravitational-wave amplitude comparisons are also done between numerical simulations and post-Newtonian, and the agreement depends on the post-Newtonian order of the amplitude expansion: the amplitude difference is about 6–7% for zeroth order and becomes smaller for increasing order. A newly derived 3.0PN amplitude correction improves agreement significantly (< 1% amplitude difference throughout most of the run, increasing to 4% near merger) over the previously known 2.5PN amplitude terms.

PACS numbers: 04.25.D-, 04.25.dg, 04.25.Nx, 04.30.-w, 04.30.Db, 02.70.Hm

I. INTRODUCTION

The last two years have witnessed tremendous progress in simulations of black hole binaries, starting with the first stable simulation of orbiting and merging black holes [1, 2], development of the moving puncture method [3, 4] and rapid progress by other groups [5–10]. Since then, an enormous amount of work has been done on the late inspiral and merger of black hole binaries, among them studies of the universality of the merger waveforms [11, 12], investigations into black hole kicks [13–26] and spin dynamics [27–29], comparisons to post-Newtonian models [30–32], and applications to gravitational wave data analysis [33–35].

Compared to the intense activity focusing on simulations close to merger, there have been relatively few simulations covering the inspiral phase. To date, only three simulations [36–40] cover more than five orbits. Long inspiral simulations are challenging for a variety of reasons: First, the orbital period increases rapidly with separation, so that simulations must cover a significantly longer evolution time. In addition, the gravitational waveform must be extracted at larger radius (and the simulation must therefore cover a larger spatial volume) because the gravitational wavelength is longer. Furthermore, gravitational wave data analysis requires small *absolute* accumulated phase uncertainties in the waveform, so the relative

phase uncertainty of the simulation must be smaller.

Gravitational wave detectors provide a major driving force for numerical relativity (NR). The first generation interferometric gravitational wave detectors, such as LIGO [41, 42], GEO600 [43] and VIRGO [44, 45], are now operating at or near their design sensitivities. Furthermore, the advanced generation of detectors are entering their construction phases. This new generation of interferometers will improve detector sensitivity by a factor of ~ 10 and hence increase expected event rates by a factor of ~ 1000 [46]. One of the most promising sources for these detectors is the inspiral and merger of binary black holes (BBHs) with masses $m_1 \sim m_2 \sim 10\text{--}20 M_\odot$ [47]. These systems are expected to have circularized long before their gravitational waves enter the sensitive frequency band of ground-based detectors [48].

A detailed and accurate understanding of the gravitational waves radiated as the black holes spiral towards each other will be crucial not only to the initial detection of such sources, but also to maximize the information that can be obtained from signals once they are observed. When the black holes are far apart, the gravitational waveform can be accurately computed using a post-Newtonian (PN) expansion. As the holes approach each other and their velocities increase, the post-Newtonian expansion is expected to diverge from the true waveform. It is important to quantify any differences

between theoretical waveforms and the true signals, as discrepancies will cause a reduction of search sensitivity. Several techniques have been proposed to address the problem of the breakdown of the post-Newtonian approximation [49–51], but ultimately, the accuracy of the post-Newtonian waveforms used in binary black hole gravitational wave searches can only be established through comparisons with full numerical simulations.

Unfortunately, comparing post-Newtonian approximations to numerical simulations is not straightforward, the most obvious problem being the difficulty of producing long and sufficiently accurate numerical simulations as explained above. In addition, post-Newtonian waveforms typically assume circular orbits, and most astrophysical binaries are expected to be on circular orbits late in their inspiral, so the orbital eccentricity within the numerical simulation must be sufficiently small¹. Another factor that complicates comparisons is the variety of post-Newtonian approximants available, from several straightforward Taylor expansions to more sophisticated Padé resummation techniques and the effective one-body approach (see e.g. [49, 50, 52–58], as well as Section III E below). While all post-Newtonian approximants of the same order should agree sufficiently early in the inspiral (when neglected higher-order terms are small), they begin to disagree with each other during the late inspiral when the post-Newtonian approximation starts to break down—exactly the regime in which NR waveforms are becoming available.

Finally, agreement (or disagreement) between NR and PN waveforms will also depend very sensitively on the precise protocol used to compare the waveforms. Are PN and NR waveforms matched early or late in the inspiral? Is the matching done at a particular time, or is a least-squares fit performed over part (or all) of the waveform? Does one compare frequencies $\omega(t)$ or phases $\phi(t)$? Are comparisons presented as functions of time or of frequency? Up to which cutoff frequency does one compare PN with NR?

Despite these difficulties, several comparisons between NR and PN have been done for the last few orbits of an equal-mass, non-spinning black hole binary. The first such study was done by Buonanno *et al* [32, 33] based on simulations performed by Pretorius [1] lasting somewhat more than 4 orbits (~ 8 gravitational wave cycles). This comparison finds visual agreement between the numerical evolution and a particular post-Newtonian approximant (in our language TaylorT3 3.0/0.0²). However, as the

authors note, this study is severely limited by numerical resolution, sizable initial eccentricity (~ 0.015), close initial separation of the black holes, and coordinate artifacts; for these reasons, the authors do not quantify the level of agreement.

More recently, Baker *et al.* [37, 38] performed simulations covering the last ~ 14 cycles before merger. These simulations have an orbital eccentricity ~ 0.008 [37], forcing the authors to compare quasi-circular post-Newtonian waveforms against a fitted smooth (“de-eccentricized”) gravitational wave phase, rather than the actual gravitational wave phase of their evolution. Comparing to TaylorT4 3.5/2.5, they find agreement between numerical and post-Newtonian gravitational wave phase to within their numerical errors, which are about 2 radians. The authors also indicate that other post-Newtonian approximants do not match their simulation as well as TaylorT4, but unfortunately, they do not mention whether any disagreement is significant (i.e., exceeding their numerical errors).

The most accurate inspiral simulation to date was performed by the Jena group and presented in Husa *et al.* [40] and Hannam *et al.* [39]. This simulation covers 18 cycles before merger and has an orbital eccentricity of ~ 0.0018 [59]. Discarding the first two cycles which are contaminated by numerical noise, and terminating the comparison at a gravitational-wave frequency $m\omega = 0.1$ (see Eq. (15) for the precise definition) their comparison extends over 13 cycles. We discuss the results of Ref. [39] in more detail in Sec. VI A 1.

This paper presents a new inspiral simulation of a non-spinning equal mass black hole binary. This new simulation more than doubles the evolution time of the simulations in Refs. [37–40], resulting in a waveform with 30 gravitational wave cycles, ending ~ 1.5 cycles before merger, and improves numerical truncation errors by one to two orders of magnitude over those in Refs. [37–40]. The orbital eccentricity of our simulations is $\sim 6 \times 10^{-5}$; this low eccentricity is achieved using refinements of techniques described in [36]. We present a detailed analysis of various effects which might influence our comparisons to post-Newtonian waveforms for non-spinning black hole binaries on circular orbits. These effects result in an uncertainty of ~ 0.05 radians out of the accumulated ~ 200 radians. Perhaps surprisingly, the largest uncertainty arises from the residual orbital eccentricity, despite its tiny value. The second largest effect arises due to a potential residual spin on the black holes, which we bound by $|S|/M_{\text{irr}}^2 < 5 \times 10^{-4}$.

We compare the numerical waveforms with three different time-domain post-Newtonian Taylor-approximants [51–53] and we match PN and NR waveforms at a specific time during the inspiral. We explore the effects of varying this matching time. When

¹ Unfortunately, this circularization occurs on extremely long time scales [48], thousands of orbits, making it impossible to run the numerical simulation long enough to radiate the eccentricity away.

² We identify post-Newtonian approximants with three pieces of information: the label introduced by [52] for how the orbital phase is evolved; the PN order to which the orbital phase is computed; and the PN order that the amplitude of the waveform

is computed. See Sec. III E for more details.

matching ~ 9 cycles after the start of our evolution, all post-Newtonian approximants of 3.0PN and 3.5PN order in orbital phase agree with our simulation to within ~ 0.03 radians over these first 9 cycles. This agreement is better than the combined uncertainties of the comparison, thus validating our simulations in a regime where the 3.5PN truncation error of post-Newtonian theory is comparable to the accuracy of our simulations. Lower order post-Newtonian approximants (2.0PN and 2.5PN order), however, accumulate a significant phase difference of ~ 0.2 radians over this region.

Extending the comparison toward merger (as well as when matching closer to merger), we find, not surprisingly, that the agreement between PN and NR at late times depends strongly on exactly what post-Newtonian approximant we use [52, 53]. Typical accumulated phase differences are on the order of radians at frequency $m\omega = 0.1$. One particular post-Newtonian approximant, TaylorT4 at 3.5PN order in phase, agrees with our NR waveforms far better than the other approximants, the agreement being within the phase uncertainty of the comparison (0.05 radians) until after the gravitational wave frequency passes $m\omega = 0.1$ (about 3.5 cycles before merger). It remains to be seen whether this agreement is fundamental or accidental, and whether it applies to more complicated situations (e.g. unequal masses, nontrivial spins).

We also compare the post-Newtonian gravitational wave amplitude to the numerical amplitude, where we estimate the uncertainty of this comparison to be about 0.5%. Restricted waveforms (i.e., 0PN order in the amplitude expansion) are found to disagree with the numerical amplitudes by 6–7%. An amplitude expansion of order 2PN shows significantly better agreement than the expansion at order 2.5PN. A newly derived 3PN amplitude [60] is found to give much better agreement than the 2.0PN amplitude.

This paper is organized as follows: Section II discusses our numerical techniques. In particular, we describe how we construct binary black hole initial data, evolve these data for 15 orbits, extract gravitational wave information from the evolution, and produce a gravitational waveform as seen by an observer at infinity. Section III details the generation of post-Newtonian waveforms, including details of how we produce the three approximants that we compare against NR. We describe our procedure for comparing NR and PN waveforms in Sec. IV, and present a detailed study of various sources of uncertainty in Sec. V. The comparisons between NR and PN are presented in Section VI. This section is split into two parts: First, we compare each PN approximant separately with the numerical simulation. Subsequently, we show some additional figures which facilitate cross-comparisons between the different PN approximants. Finally, we present some concluding remarks in Section VII. The impatient reader primarily interested in NR-PN comparisons may wish to proceed directly to Table III summarizing the uncertainties of our comparisons, and then continue to Sec. VI,

starting with Fig. 15.

II. GENERATION OF NUMERICAL WAVEFORMS

In order to do a quantitative comparison between numerical and post-Newtonian waveforms, it is important to have a code capable of starting the black holes far enough apart to be in a regime where we strongly believe the post-Newtonian approximation is valid, track the orbital phase extremely accurately, and do so efficiently so the simulation can be completed in a reasonable amount of time. Furthermore, the gravitational waves from such a simulation must be extracted in such a manner that preserves the accuracy of the simulation and predicts the waveform as seen by a distant observer, so a comparison with the post-Newtonian waveform can be made. In this section we describe the techniques we use to do this, as well as the results of a simulation starting more than 15 orbits prior to merger.

When discussing numerical solutions of Einstein's equations, we write all dimensionful quantities in terms of some mass scale m , which we choose to be the sum of the irreducible masses of the two black holes in the initial data:

$$m = M_{\text{irr},1} + M_{\text{irr},2}. \quad (1)$$

The irreducible mass of a single hole is defined as

$$M_{\text{irr}} \equiv \sqrt{A/16\pi}, \quad (2)$$

where A is the surface area of the event horizon; in practice we take A to be the surface area of the apparent horizon. More generally, it is more appropriate to use the Christoudoulou mass of each black hole,

$$M_{\text{BH}}^2 = M_{\text{irr}}^2 + \frac{S^2}{4M_{\text{irr}}^2}, \quad (3)$$

instead of the irreducible mass. Here S is the spin of the hole. However, for the case considered in this paper, the spins are sufficiently small that there is little difference between M_{BH} and M_{irr} .

A. Initial data

Initial data are constructed within the conformal thin sandwich formalism [61, 62] using a pseudo-spectral elliptic solver [63]. We employ quasi-equilibrium boundary conditions [64, 65] on spherical excision boundaries, choose conformal flatness and maximal slicing, and use Eq. (33a) of Ref. [66] as the lapse boundary condition. The spins of the black holes are made very small via an appropriate choice of the tangential shift at the excision surfaces, as described in [66].

As the most accurate post-Newtonian waveforms available assume adiabatic inspiral of quasi-circular orbits, it

TABLE I: Summary of the initial data sets used in this paper. The first block of numbers (d , Ω_0 , f_r , and v_r) represent raw parameters entering the construction of the initial data. The second block gives some properties of each initial data set: m denotes the sum of the irreducible masses, M_{ADM} and J_{ADM} the ADM energy and angular momentum, and s_0 the initial proper separation between the horizons. The last column lists the eccentricity computed from Eq. (7). The initial data set 30c is used for all evolutions (except for consistency checks) described in this paper.

Name	d	Ω_0	f_r	$v_r \times 10^4$	$m\Omega_0$	M_{ADM}/m	J_{ADM}/m^2	s_0/m	$e_{ds/dt}$
30a	30	0.0080108	0.939561	0.00	0.01664793	0.992333	1.0857	17.37	1.0×10^{-2}
30b	30	0.0080389	0.939561	-4.90	0.0167054	0.992400	1.0897	17.37	6.5×10^{-4}
30c	30	0.0080401	0.939561	-4.26	0.0167081	0.992402	1.0898	17.37	5×10^{-5}
24a	24	0.0110496	0.92373	-8.29	0.0231947	0.990759	1.0045	14.15	1.1×10^{-3}
24b	24	0.0110506	0.923739	-8.44	0.0231967	0.990767	1.0049	14.15	1.5×10^{-4}




FIG. 1: Proper separation (top panel) and its time derivative (lower panel) versus time for short evolutions of the $d = 30$ initial data sets '30a', '30b', and '30c' (see Table I). These three data sets represent zero through two iterations of our eccentricity-reduction procedure. The orbital eccentricity is reduced significantly by each iteration.

is desirable to reduce the eccentricity of the numerical data as much as possible. Using techniques developed in [36], each black hole is allowed to have a nonzero initial velocity component towards the other hole. This small velocity component v_r and the initial orbital angular velocity Ω_0 are then fine-tuned in order to produce an orbit with very small orbital eccentricity³. We have improved our eccentricity-reduction procedure since the version described in [36], so we summarize our new iterative procedure here:

³ An alternative method of producing low-eccentricity initial data, based on post-Newtonian ideas, is developed in [59]. While that technique is computationally more efficient than ours, it merely reduces orbital eccentricity by a factor of ~ 5 relative to quasi-circular initial data, which is insufficient for the comparisons presented here.

We start with a quasi-circular (i.e., $v_r = 0$) initial data set at coordinate separation $d = 30$, where Ω_0 is determined by equating Komar mass with ADM mass [66]. We then evolve these data for about 1.5 orbits, corresponding to a time $t/m \approx 600$. From this short evolution, we measure the time derivative of the proper separation between the horizons, ds/dt (which is plotted in Fig. 1). We fit ds/dt in the interval $100 \lesssim t/m \lesssim 600$ to the function

$$\frac{ds}{dt} = A_0 + A_1 t + B \cos(\omega t + \varphi), \quad (4)$$

where we vary all five parameters A_0 , A_1 , B , ω and φ to achieve the best fit. The desired smooth inspiral is represented by the part $A_0 + A_1 t$; the term $B \cos(\omega t + \varphi)$ corresponds to oscillations caused by orbital eccentricity. We then choose new values of v_r and Ω_0 such that a *Newtonian* orbit with radial velocity $B \cos(\omega t + \varphi)$ at initial separation s_0 would be perfectly circular. This is accomplished by the updating formulae

$$\Omega_0 \rightarrow \Omega_0 + \frac{B \sin \varphi}{2s_0}, \quad (5)$$

$$v_r \rightarrow v_r - \frac{B \cos \varphi}{2}, \quad (6)$$

where s_0 is the initial proper separation between the horizons. We then use the new values of Ω_0 and v_r to construct a new initial data set, we again evolve for two orbits, fit to Eq. (4), and again update Ω_0 and v_r . We continue iterating this procedure until the eccentricity is sufficiently small.

We estimate the eccentricity for each iteration from the fit to Eq. (4) using the formula

$$e_{ds/dt} = \frac{B}{s_0 \omega}, \quad (7)$$

which is valid in Newtonian gravity for small eccentricities. Successive iterations of this procedure yield the initial data sets 30a, 30b, and 30c summarized in Table I. Eccentricity decreases by roughly a factor of 10 in each iteration, with 30c having $e_{ds/dt} \approx 5 \times 10^{-5}$. The evolutions used during eccentricity reduction need not be very accurate and need to run only for a short time, $t \sim 600m$. One iteration of this procedure at our

second lowest resolution requires about 250 CPU-hours. For completeness, Table I also lists parameters for initial data at smaller separation; these data will be used for consistency checks below. Apart from these consistency checks, the remainder of this paper will focus exclusively on evolutions of the low-eccentricity initial data set 30c.

B. Evolution of the inspiral phase

The Einstein evolution equations are solved with the pseudo-spectral evolution code described in Ref. [10]. This code evolves a first-order representation [67] of the generalized harmonic system [68–70]. We handle the singularities by excising the black hole interiors from our grid. Our outer boundary conditions [67, 71, 72] are designed to prevent the influx of unphysical constraint violations [73–79] and undesired incoming gravitational radiation [80], while allowing the outgoing gravitational radiation to pass freely through the boundary.

The code uses a fairly complicated domain decomposition to achieve maximum efficiency. Each black hole is surrounded by several (typically six) concentric spherical shells, with the inner boundary of the innermost shell (the excision boundary) just inside the horizon. A structure of touching cylinders (typically 34 of them) surrounds these shells, with axes along the line between the two black holes. The outermost shell around each black hole overlaps the cylinders. The outermost cylinders overlap a set of outer spherical shells, centered at the origin, which extend to large outer radius. External boundary conditions are imposed only on the outer surface of the largest outer spherical shell. We vary the location of the outer boundary by adding more shells at the outer edge. Since all outer shells have the same angular resolution, the cost of placing the outer boundary farther away (at full resolution) increases only linearly with the radius of the boundary. External boundary conditions are enforced using the method of Bjorhus [81], while inter-domain boundary conditions are enforced with a penalty method [82, 83].

We employ the dual-frame method described in Ref. [10]: we solve the equations in an ‘inertial frame’ that is asymptotically Minkowski, but our domain decomposition is fixed in a ‘comoving frame’ that rotates with respect to the inertial frame and also shrinks with respect to the inertial frame as the holes approach each other. The positions of the holes are fixed in the comoving frame; we account for the motion of the holes by dynamically adjusting the coordinate mapping between the two frames. Note that the comoving frame is referenced only internally in the code as a means of treating moving holes with a fixed domain. Therefore all coordinate quantities (e.g. black hole trajectories, wave-extraction radii) mentioned in this paper are inertial-frame values unless explicitly stated otherwise.

One side effect of our dual frame system is that the outer boundary of our domain (which is fixed in the co-

moving frame) moves inward with time as observed in the inertial frame. This is because the comoving frame shrinks with respect to the inertial frame to follow the motion of the holes. In Refs. [10, 36] the inertial frame coordinate radius r (with respect to the center of mass) and the comoving coordinate radius r' are related by a simple scaling

$$r = a(t)r'. \quad (8)$$

The expansion parameter $a(t)$ is initially set to unity and decreases dynamically as the holes approach each other, so that the comoving-frame coordinate distance between the holes remains constant. The outer boundary of the computational grid is at a fixed comoving radius R'_{bdry} , which is mapped to the inertial coordinate radius $R_{\text{bdry}}(t) = a(t)R'_{\text{bdry}}$. Because we wish to accurately compute the gravitational radiation as measured far from the holes, it is desirable to have a moderately large outer boundary ($R_{\text{bdry}}(t) \gtrsim 200m$) throughout the run. For the linear mapping, Eq. (8), this requires a very distant outer boundary early in the run, $R_{\text{bdry}}(0) \simeq 1000m$. Computationally this is not very expensive. However, the initial junk radiation contaminates the evolutions for a time interval proportional to the light-crossing time to the outer boundary, and for $R_{\text{bdry}}(0) \simeq 1000m$ it would be necessary to discard a significant portion of the evolution.

We therefore use the mapping

$$r = \left[a(t) + (1 - a(t)) \frac{r'^2}{R_0^2} \right] r', \quad (9)$$

for some constant R'_0 which is chosen to be roughly the radius of the outer boundary in comoving coordinates. This mapping has the following properties: (1) At the initial time $t = 0$, the map reduces to the identity map because $a(0) = 1$. Thus we do not need to re-map our initial data before evolving. (2) For small radii (i.e., at the locations of the black holes), the map reduces to the linear map, $r = a(t)r' + \mathcal{O}(r'^3)$. This allows use of the control system without modifications. (3) The moving radius $r' = R'_0$ is mapped to a *constant* inertial radius: $r(R'_0) = R'_0$. This allows us to keep the inertial radius of the outer boundary constant (or nearly constant⁴) in time rather than shrinking rapidly.

In total, we have run three evolutions of the 30c initial data set; these use different combinations of outer boundary radius and radial mapping between inertial and moving coordinates. Some properties of these evolutions are summarized in Table II. We also performed extensive convergence testing, running the same evolution on

⁴ In practice, we choose R'_0 somewhat larger than the outer boundary, so that the outer boundary of the computational domain slowly contracts in inertial coordinates. This makes the zero-speed characteristic fields *outgoing* there, avoiding the need to impose boundary conditions on those fields.

TABLE II: Overview of low-eccentricity simulations discussed in this paper. R_{bdry} is the initial coordinate radius of the outer boundary; this radius changes during the evolution according to the choice of “radial map” between inertial and comoving coordinates. The last column lists the different resolutions run for each evolution, N6 being highest resolution. Evolution 30c-1/N6 forms the basis of our post-Newtonian comparisons, and is used in all figures unless noted otherwise.

Name	ID	N_{orbits}	R_{bdry}	radial map	resolutions
30c-1	30c	15.6	462m	Eq. (9)	N1, N2, . . . , N6
30c-2	30c	15.6	722m	Eq. (8)	N2, N4, N6
30c-3	30c	15.6	202m	Eq. (8)	N2, N3, . . . , N6
24b-1	24b	8.3	160m	Eq. (8)	N2, N3, N4




FIG. 2: Spacetime diagram showing the spacetime volume simulated by the numerical evolutions listed in Tab. II. The magnified view in the right panel shows how the gravitational waves are escorted to our extraction radii (see Sec. II C) after the simulation in the center has already crashed at $t \sim 3930m$, and after the estimated time of the black hole merger, which is indicated by the circle. The thin diagonal lines are lines of constant $t - r^*$; each corresponds to a retarded time at which the gravitational wave frequency ω at infinity assumes a particular value.

up to six distinct resolutions, N1 to N6. The coarsest resolution 30c-1/N1 uses approximately 41^3 grid points (summing all grid points in all the subdomains), while the most accurate evolution, 30c-1/N6, uses about 67^3 grid points. The run 30c-1/N2 required about 2,500 CPU-hours and run 30c-1/N6 about 19,000, where our simulations do not take advantage of symmetries. The distance to the outer boundary is adjusted by adding or removing outer spherical shells to an otherwise unmodified domain-decomposition. Run 30c-1 has 20 such outer spherical shells, while 30c-2 utilizes 32 and 30c-3 only 8. Thus, the total number of grid points varies slightly between runs, e.g. about 71^3 for 30c-2/N6. Figure 2 indicates the dif-




FIG. 3: Coordinate trajectories of the centers of the black holes. The small circles/ellipsoids show the apparent horizons at the initial time and at the time when the simulation ends and wave escorting begins. The inset shows an enlargement of the dashed box.

ferent behavior of the outer boundary location for these three evolutions.

For all of the evolutions 30c-1/2/3, the coordinate trajectories of the centers of the apparent horizons appear as in Fig. 3. The regular inspiral pattern without noticeable oscillations once again indicates that our evolutions indeed have very low eccentricity.

Figure 4 demonstrates the convergence of the black hole mass $m(t)$ with spatial resolution for run 30c-1. The mass $m(t)$ is computed as the sum of the irreducible masses of both black holes, as defined in Eq. (2). At the highest resolution, $m(t)$ deviates by only a few parts in 10^6 from its initial value m .

We also measure the quasi-local spin using coordinate rotation vectors projected into the apparent horizon surfaces [84–86]. Only the z-component of the spin is non-zero (i.e., the spins are aligned with the orbital angular momentum). The spin starts at $S_z/M_{\text{irr}}^2 \approx -6 \times 10^{-5}$ and increases slowly to -5×10^{-4} during the evolution, where the minus sign indicates that the black hole spin is anti-aligned with the orbital angular momentum. Thus it appears the black hole’s spins move further away from the corotational state. We believe this effect is caused by the use of coordinate rotation vectors when calculating the quasi-local spin, rather than more sophisticated approximate Killing vectors [87–89]. Preliminary results with approximate Killing vectors find the initial spin to be less than 10^{-6} , and slowly increasing during the evolution to a final value of 2×10^{-5} at the end of the comparison interval to post-Newtonian theory. Given the preliminary character of these results, we will take here the conserva-




FIG. 4: Deviation of total irreducible mass $m(t) = 2M_{\text{irr}}(t)$ from its value in the initial data. Plotted are the five different resolutions of run 30a-1.

tive bound $|\mathbf{S}|/M_{\text{irr}}^2 \leq 5 \times 10^{-4}$ obtained from coordinate rotation vectors.

C. Escorting gravitational waves

The simulation presented in Fig. 3 stops when the horizons of the black holes become too distorted just before merger. At that point, the proper separation between the horizons is $\sim 4.0m$, and the orbital frequency has reached $m\Omega_{\text{orbit}} = 0.125$; comparison with [32] suggests this is about $15m$ before formation of a common apparent horizon. So far our efforts to simulate the actual merger and ringdown have been unsuccessful.

At the time the simulation ends, most of the domain (all regions except for the immediate vicinity of the two holes) is still well resolved, and the spacetime contains gravitational radiation that has not yet propagated out to the large radii where we perform wave extraction. So instead of losing this information, which consists of several gravitational-wave cycles, we evolve only the outer portions of our grid beyond the time at which the code crashes in the center, effectively ‘escorting’ the radiation out to the extraction radii.

To do this, we first stop the evolution shortly before it crashes, and we introduce a new spherical excision boundary that surrounds both black holes and has a radius of roughly three times the black hole separation. This new excision boundary moves radially outward at slightly faster than the speed of light so that it is causally disconnected from the interior region where the code is crashing, and so that no boundary conditions are required on this boundary. We then continue the evolution on the truncated spherical-shell domain that extends from the new excision boundary to the outer boundary. To move both boundaries appropriately, we employ a new radial coordinate mapping

$$r = A(t)r(r') + B(t), \quad (10)$$




FIG. 5: Gravitational waveform extracted at $r = 240m$. From top panel to bottom: The real part of the $l = 2, m = 2$ component of $r\Psi_4$; the gravitational wave strain, obtained by two time integrals of $\text{Re}(r\Psi_4)$; the frequency of the gravitational wave, Eq. (15); the gravitational wavelength, $\lambda = 2\pi/\omega$. The vertical grey line at $t \approx 3930m$ indicates the time when ‘wave escorting’ starts.

where $r(r')$ is given by Eq. (9). The functions $A(t)$ and $B(t)$ are chosen to satisfy three criteria: First, the inner boundary of the spherical shell moves outward with coordinate speed of unity, which turns out to be slightly superluminal. Second, the outer boundary location $R_{\text{bdry}}(t)$ has continuous first and second time derivatives at the time we transition to the truncated domain. And finally, the outer boundary location $R_{\text{bdry}}(t)$ approaches some fixed value at late times. The right panel of Fig. 2 shows the motion of the inner and outer radii for evolutions 30c-1 and 30c-2 (we did not perform wave escorting for 30c-3). For 30c-1, wave escorting extends the evolution for an additional time $220m$ beyond the point at which the simulation stops in the center.

Figure 5 shows the gravitational waveform extracted at inertial coordinate radius $R = 240m$ for the run 30c-1. The grey vertical line indicates the time when wave escorting starts. Wave escorting allows us to extract another 4 cycles of gravitational waves. The gravitational wave frequency at the end of the run is $m\omega \approx 0.16$, placing the end of the waveform at about $50m$ (or ~ 1.5 cycles) before formation of a common apparent horizon⁵ (judged by comparison with [32]).

⁵ The waveform ends somewhat further from merger than the orbital trajectory, because the artificial boundary is placed initially at a radius $\sim 15m$, and then moves outward somewhat faster than the speed of light, thus overtaking the very last part of the waveform as it travels to the wave-extraction radii.




FIG. 6: Constraint violations of run 30c-1. The top panel shows the L^2 norm of all constraints, normalized by the L^2 norm of the spatial gradients of all dynamical fields. The bottom panel shows the same data, but without the normalization factor. Norms are taken only in the regions outside apparent horizons.

In the lower two panels of Fig. 5 there is a significant amount of noise near the beginning of the run, at $t < 250m$. This noise is barely evident in the top panel of Fig. 5 as well. The noise is a manifestation of ‘junk radiation’, a pulse of radiation often seen at the beginning of numerical relativity simulations, and is caused by the initial data not being precisely a snapshot of an evolution that has been running for a long time. Among the effects that produce junk radiation are incorrect initial distortions of the individual holes, so that each hole radiates as it relaxes to its correct quasi-equilibrium shape.

Our evolution code does not explicitly enforce either the Einstein constraints or the secondary constraints that arise from writing the system in first-order form. Therefore, examining how well these constraints are satisfied provides a useful consistency check. Figure 6 shows the constraint violations for run 30c-1. The top panel shows the L^2 norm of all the constraint fields of our first order generalized harmonic system, normalized by the L^2 norm of the spatial gradients of the dynamical fields (see Eq. (71) of Ref. [67]). The bottom panel shows the same quantity, but without the normalization factor (*i.e.*, just the numerator of Eq. (71) of Ref. [67]). The L^2 norms are taken over the entire computational volume that lies outside of apparent horizons. At early times, $t < 500m$, the constraints converge rather slowly with resolution because the junk radiation contains high frequencies. Convergence is more rapid during the smooth inspiral phase, after the junk radiation has exited through the outer boundary. The constraints increase around $t \sim 3900m$ as the code begins to fail near the two merging holes, but then the constraints decrease again after the failing

region is excised for wave escorting. The normalized constraint violations are less than 10^{-4} until just before the peak (which occurs at $t = 3930m$ for all but the lowest resolutions). The size of the peak causes some concern that the waveforms at late times may be contaminated by constraint violations to a non-negligible degree. However, near the peak, the constraint violations are large only in the inner regions of the domain near the black holes (note that the curves in Fig. 6 decrease by two orders of magnitude immediately after these inner regions are excised at $t = 3930m$). Because all constraint quantities propagate at the speed of light or slower for the formulation of Einstein’s equations that we use, any influence that the constraint peak has on the extracted waveform occurs after the constraint violations have had time to propagate out to the wave extraction zone. This is very late in the waveform, well after the gravitational wave frequency reaches $m\omega = 0.1$, as can be seen from the right panel of the spacetime diagram in Fig. 2.

D. Waveform extraction

Gravitational waves are extracted using the Newman-Penrose scalar Ψ_4 , using the same procedure as in [36]. To summarize, given a spatial hypersurface with timelike unit normal n^μ , and given a spatial unit vector r^μ in the direction of wave propagation, the standard definition of Ψ_4 is the following component of the Weyl curvature tensor,

$$\Psi_4 = -C_{\alpha\mu\beta\nu}\ell^\mu\ell^\nu\bar{m}^\alpha\bar{m}^\beta, \quad (11)$$

where $\ell^\mu \equiv \frac{1}{\sqrt{2}}(n^\mu - r^\mu)$, and m^μ is a complex null vector (satisfying $m^\mu\bar{m}_\mu = 1$) that is orthogonal to r^μ and n^μ . Here an overbar denotes complex conjugation.

For (perturbations of) flat spacetime, Ψ_4 is typically evaluated on coordinate spheres, and in this case the usual choices for n^μ , r^μ and m^μ are

$$n^\mu = \left(\frac{\partial}{\partial t}\right)^\mu, \quad (12a)$$

$$r^\mu = \left(\frac{\partial}{\partial r}\right)^\mu, \quad (12b)$$

$$m^\mu = \frac{1}{\sqrt{2}r} \left(\frac{\partial}{\partial \theta} + i\frac{1}{\sin\theta}\frac{\partial}{\partial \phi}\right)^\mu, \quad (12c)$$

where (r, θ, ϕ) denote the standard spherical coordinates. With this choice, Ψ_4 can be expanded in terms of spin-weighted spherical harmonics of weight -2 :

$$\Psi_4(t, r, \theta, \phi) = \sum_{lm} \Psi_4^{lm}(t, r) {}_{-2}Y_{lm}(\theta, \phi), \quad (13)$$

where the Ψ_4^{lm} are expansion coefficients defined by this equation.

For curved spacetime, there is considerable freedom in the choice of the vectors r^μ and m^μ , and different researchers have made different choices [5, 32, 90–94] that

are all equivalent in the $r \rightarrow \infty$ limit. We choose these vectors by first picking an extraction two-surface \mathcal{E} that is a coordinate sphere ($r^2 = x^2 + y^2 + z^2$) centered on the center of mass of the binary system (using the global asymptotically Cartesian coordinates employed in our code). We choose r^μ to be the outward-pointing spatial unit normal to \mathcal{E} (that is, we choose r_i proportional to $\nabla_i r$ and raise the index with the spatial metric). Then we choose m^μ according to Eq. (12c), using the standard spherical coordinates θ and ϕ defined on these coordinate spheres. Finally we use Eqs. (11) and (13) to define the Ψ_4^{lm} coefficients.

Note that the m^μ vector used here is not exactly null nor exactly of unit magnitude at finite r . The resulting Ψ_4^{lm} at finite r will disagree with the waveforms observed at infinity. Our definition does, however, agree with the standard definition given in Eqs. (11)–(13) as $r \rightarrow \infty$. Because we extrapolate the extracted waves to find the asymptotic radiation field (see Section II F), these effects should not play a role in our PN comparisons: Relative errors in Ψ_4^{lm} introduced by using the simple coordinate tetrad fall off like $1/r$, and thus should vanish after extrapolating to obtain the asymptotic behavior. While more careful treatment of the extraction method—such as those discussed in [95–97]—may improve the quality of extrapolation and would be interesting to explore in the future, the naive choice made here should be sufficient to ensure that the waveform after extrapolation is correct to the accuracy needed for these simulations.

In this paper, we focus on the $l = 2, m = 2$ coefficient. Following common practice (see e.g. [5, 11]), we split the extracted waveform into real phase ϕ and real amplitude A , defined by

$$\Psi_4^{22}(r, t) = A(r, t)e^{-i\phi(r, t)}. \quad (14)$$

The gravitational-wave frequency is given by

$$\omega = \frac{d\phi}{dt} \quad (15)$$

The minus sign in the definition of ϕ is chosen so that the phase increases in time and ω is positive. Equation (14) defines ϕ only up to multiples of 2π . These multiples of 2π are chosen to make ϕ continuous through each evolution, still leaving an overall multiple of 2π undetermined. We will consider only phase differences in this paper, so the choice of this overall phase offset is irrelevant.

E. Convergence of extracted waveforms

In this section we examine the convergence of the gravitational waveforms extracted at fixed radius, without extrapolation to infinity. This allows us to study the behavior of our code without the complications of extrapolation. The extrapolation process and the resulting extrapolated waveforms are discussed in Sec. II F.

The top panel of Fig. 7 shows the convergence of the gravitational wave phase ϕ with numerical resolution for




FIG. 7: Convergence of the gravitational wave phase extracted at radius $R = 77m$. All lines show differences with respect to our highest resolution run, 30c-1/N6. The top panel shows different resolutions of the same run 30c-1; no time or phase shifts have been performed. The bottom panel compares different runs, aligning the runs at $m\omega = 0.1$ by a time and phase shift. The thin vertical line indicates the time at which $m\omega = 0.1$ for 30c-1/N6.

the run 30c-1. For this plot, the waveform is extracted at a fixed radius $R = 77m$. Each line shows the difference between ϕ computed at some particular resolution and ϕ computed from our highest-resolution run 30c-1/N6. When subtracting results at different resolutions, no time or phase adjustment has been performed. The difference in ϕ between the two highest-resolution runs is smaller than 0.03 radians throughout the run, and it is smaller than 0.02 radians between $t = 1000m$ and the point at which $m\omega = 0.1$.

At times before $1000m$, the phase convergence of our simulation is limited to about 0.05 radians because of effects of junk radiation (described at the end of Section II C). The sharp pulse of junk radiation has comparatively large numerical truncation error, and excites all characteristic modes at truncation-error level, including waves that propagate back toward the origin. Generation of these secondary waves stops when the pulse of junk radiation leaves through the outer boundary (*i.e.*, after one light-crossing time). Because we use the improved outer boundary conditions of Rinne *et al.* [72], there are no significant reflections when the junk radiation passes through the outer boundary. However, the waves produced before the junk radiation leaves remain in the computational domain for two additional light-crossing times, until they eventually leave through the outer boundary.

The bottom panel of Fig. 7 shows phase comparisons between different waveforms after we perform a time shift and phase shift so that the waveforms agree at $m\omega = 0.1$.

Our procedure for time shifting and phase shifting is the same as the shifting procedure we use to compare NR with PN waveforms (see Sec. IV B), so that the error estimates we extract from the bottom panel of Fig. 7 are relevant for our later NR-PN comparison.

There are three different types of comparisons shown in the bottom panel of Fig. 7: Phase differences between runs with the same initial data but with different outer boundary locations, phase differences between runs with different initial data, and phase differences between different numerical resolutions of the same run (this last comparison is the same as what is shown in the top panel, except in the bottom panel the waveforms are time and phase shifted). We will discuss all three of these in turn.

First, we compare the phase difference of 30c-1/N6 with runs that have different outer boundary locations. Run 30c-2 (with more distant outer boundary) agrees to within 0.002 radians with run 30c-1, but run 30c-3 (with closer outer boundary), has a much larger phase difference with 30c-1. We believe that this is because run 30c-3 has a very small ratio of outer boundary location to gravitational wavelength: R/λ is about 1.1 for the first two-thirds of the run, and remains less than 2 for the entire run.

We can explain the order of magnitude of these phase differences using the analysis of Buchman & Sarbach [80]. Our outer boundary conditions are not perfectly absorbing, but instead they reflect some fraction of the outgoing radiation.⁶ The ratio of the amplitude of curvature perturbations (i.e. Ψ_4) of the reflected wave to that of the outgoing wave is

$$q \approx \frac{3}{2(2\pi)^4} \left(\frac{\lambda}{R} \right)^4. \quad (16)$$

Although this fraction is computed at the outer boundary, the ratio of reflected to outgoing amplitudes is also approximately q near the black holes because the incoming reflected waves grow like $1/r$ as they travel inward just like the outgoing waves decrease by $1/r$ as they propagate outward. Now consider the second time derivative of the gravitational wave phase, $\ddot{\phi}$; this is nonzero only because of gravitational wave emission, so $\ddot{\phi}$ is proportional to some power of the outgoing wave amplitude. To get the correct power, we can use Eq. (45) to find $\dot{x} \sim x^5$, so Eq. (36) yields $\ddot{\phi} \sim x^{11/2}$ (we assume gravitational wave phase is twice the orbital phase). The amplitude of Ψ_4 scales like x^4 , so $\ddot{\phi} \sim A^{11/8}$. Let us assume for the sake of this rough error estimate that the change in $\ddot{\phi}$ due to the *ingoing* reflected wave scales similarly with amplitude, $\ddot{\phi} \sim \bar{A}^{11/8}$, where $\bar{A} = qA$ is the amplitude of the reflected ingoing wave. Therefore the unphysical GW

force acting back on the system due to boundary reflections will cause fractional errors in the second derivative of the phase of about $q^{11/8}$. That is, the error $\delta\phi$ caused by the improper boundary condition will be given by

$$\frac{d^2\delta\phi}{dt^2} = q^{11/8} \frac{d^2\phi}{dt^2}. \quad (17)$$

Integrating this yields $\delta\phi = q^{11/8}\phi$, where ϕ is the total gravitational wave phase accumulated during the evolution. For 30c-3, $\lambda/R \sim 0.9$, so $q \sim 6 \times 10^{-4}$, which yields $\delta\phi \sim 0.08$ radians for an accumulated gravitational wave phase of about 200 radians. This rough estimate agrees in order of magnitude with the phase difference between 30c-3 and 30c-1 as shown in the bottom panel of Figure 7. The run 30c-1 has an outer boundary about 2.5 farther away, reducing the reflection coefficient by a factor $2.5^4 \approx 40$, so for 30c-1 this estimate of the phase error gives $\delta\phi = 5 \times 10^{-4}$ radians. Therefore, we expect reflection of the outgoing radiation at the outer boundary to be insignificant for 30c-1. This is confirmed by the excellent agreement between runs 30c-1 and 30c-2 (the latter having even larger outer boundary).

The second comparison shown in the lower panel of Fig. 7 is the phase difference between 30c-1/N6 and 24b-1/N4, a shorter 8-orbit evolution started from a separate initial data set (set 24b in Table I) with a separate eccentricity-reduction procedure. The phase agreement between these two runs (including an overall time shift and phase shift) is better than 0.01 radians for a total accumulated phase of ~ 100 radians of the 8-orbit run, i.e. better than one part in 10^4 . Run 24b-1 has a similar outer boundary location as run 30c-3, and indeed both of these runs show similar phase differences from 30c-1.

Finally, the third comparison shown in the lower panel of Fig. 7 is the the phase difference between the two highest resolutions of the run 30c-1 when a time shift is applied. For $t \gtrsim 1000m$ the agreement is much better than without the time shift (see upper panel), indicating that the dominant error is a small difference in the overall evolution time. For the post-Newtonian comparisons we perform in the second part of this paper, waveforms are always aligned at specific frequencies by applying time and phase shifts. Therefore, the time-shifted phase difference as displayed in the lower panel is the most appropriate measure of numerical truncation error for these PN comparisons. This difference is less than 0.003 radians after $t = 1000m$ but is larger, about 0.02 radians, at early times where the waveforms are noisy because of junk radiation.

We now compare the gravitational wave amplitudes of different runs in the same manner as we compared the gravitational wave phases. Figure 8 presents convergence data for the amplitude of the gravitational waves for the same runs as shown in Fig. 7. Spatial truncation error for the amplitude is less than 0.1 percent for $t/m > 1000$, and earlier than this it is limited by residual noise from the junk radiation. Differences (including time shifts) between runs of different lengths are shown in the lower

⁶ However, in a comparison of various boundary conditions [72], the boundary conditions we use produced smaller reflections than other boundary conditions commonly used in numerical relativity.




FIG. 8: Convergence of the gravitational wave amplitude extracted at radius $R = 77m$. This plot corresponds to Fig. (7), except that relative amplitude differences are shown. The thin vertical line indicates the time at which $m\omega = 0.1$ for 30c-1/N6.

panel of Fig. 8. These differences are even smaller, but because of their small size, they are dominated by noise for about the first half of the run. The oscillations apparent in the comparison to 24b-1 are caused by the larger orbital eccentricity of 24b-1 (cf. Tab. I).

F. Extrapolation to infinity

The quantity of interest to gravitational wave detectors is the gravitational waveform as seen by an observer effectively infinitely far from the source. Our numerical simulations, in contrast, cover only a region of finite volume around the source, and our numerical waveforms are extracted at a finite radius. Waveforms extracted at a finite radius can differ from those extracted at infinity because of effects discussed in Section IID; these effects can lead to phase errors of several tenths of a radian and relative amplitude errors of several percent. To avoid such errors we extrapolate to infinite extraction radius as follows.

We extract data for Ψ_4 on coordinate spheres of coordinate radii $r/m = 75, 80, 85, \dots, 240$, as described in Section IID. These extracted waveforms are shifted in time relative to one another because of the finite light-travel time between these extraction surfaces. We correct for this by shifting each waveform by the tortoise-coordinate radius at that extraction point [90]

$$r^* = r_{\text{areal}} + 2M_{\text{ADM}} \ln \left(\frac{r_{\text{areal}}}{2M_{\text{ADM}}} - 1 \right). \quad (18)$$

Here M_{ADM} is the ADM mass of the initial data, and




FIG. 9: Difference between areal radius r_{areal} and coordinate radius of select extraction surfaces. r_{areal} remains constant to within $0.01m$ during the evolution. The diamond indicates M_{ADM}/m of the initial data.

$r_{\text{areal}} = \sqrt{A/4\pi}$, where A is the area of the extraction sphere. This is not the only possible choice for the retarded time—for example, the waveforms could be shifted so that the maxima of the amplitude align [39]. It has also been suggested [98] that the time shift should change with the amount of radiated energy—essentially, that the factor of M_{ADM} should be replaced by the amount of mass interior to the extraction radius at each time. We leave investigation of other choices of retarded time for future work.

Figure 9 presents the areal radius during the evolution at several typical extraction radii. The areal radius of these extraction surfaces is constant to within about $0.01m$, and to the same precision, $r_{\text{areal}} = r + M_{\text{ADM}}$. This relationship is not surprising, because the initial data is conformally flat, so that for coordinate spheres $r_{\text{areal}} = r + M_{\text{ADM}} + \mathcal{O}(M_{\text{ADM}}/r)$. For convenience, we simply set $r_{\text{areal}} = r + M_{\text{ADM}}$ in Eq. (18), rather than explicitly integrating to find the area of each extraction sphere.

After the time shift, each waveform is a function of retarded time, $t - r^*$. At a given value of retarded time, we have a series of data points—one for each extraction radius. We fit phase and amplitude of these data separately to a polynomial in $1/r$,

$$\phi(t - r^*, r) = \phi_{(0)}(t - r^*) + \sum_{k=1}^n \frac{\phi_{(k)}(t - r^*)}{r^k}, \quad (19)$$

$$rA(t - r^*, r) = A_{(0)}(t - r^*) + \sum_{k=1}^n \frac{A_{(k)}(t - r^*)}{r^k}. \quad (20)$$

The leading-order term of each polynomial, as a function of retarded time, is then the desired asymptotic waveform:

$$\phi(t - r^*) = \phi_{(0)}(t - r^*), \quad (21)$$

$$rA(t - r^*) = A_{(0)}(t - r^*). \quad (22)$$

We find good convergence of this method as we increase the order n of the extrapolating polynomial. Figure 10




FIG. 10: Error of phase extrapolation to infinity for extrapolation of order n , cf. Eq. (19). Plotted are absolute differences between extrapolation with order n and $n+1$. Increasing the order of the polynomial increases accuracy, but also amplifies noise.

shows the difference in phase between waveforms extrapolated using successively higher-order polynomials. We see a broad improvement in the accuracy of the phase with increasing order, but unfortunately, higher order extrapolations tend to amplify the noise. Our preferred choice is $n = 3$ extrapolation, resulting in extrapolation errors of $\lesssim 0.003$ radians for $t - r^* \gtrsim 1000m$.

Figure 11 is analogous to Fig. 10, except that it shows relative differences in the extrapolated amplitudes. The basic picture agrees with the phase extrapolation: Higher order extrapolation reduces the errors, but amplifies noise. Our preferred choice $n = 3$ gives a relative amplitude error of $\lesssim 0.002$ for $t - r^* \gtrsim 1000m$, dropping to less than 0.001 for $t - r^* \gtrsim 2000m$.

Phase and amplitude extrapolation become increasingly more accurate at late times. The main obstacle to accuracy seems to be near-zone effects scaling with powers of (λ/r) , where λ is the wavelength of the gravitational wave. The wavelength is quite large at the beginning of the simulation ($\lambda \approx 180m$, cf. Fig. 5), but becomes shorter during the evolution, so that even low-order extrapolation is quite accurate at late times. Alternatively, near-zone effects can be mitigated by using data extracted at large values of r . It is precisely because of these near-zone effects that we have chosen to ignore data extracted at $r < 75m$ when we extrapolate to infinity.

In Figs. 12 and 13, we show the effects of extrapolation using different ranges of extracted data. Using data extracted every $5m$ in the range $r = 50m - 90m$ results in noticeable differences early in the run—though it is adequate later in the run. For ranges at higher radii (e.g. $[75m, 150m]$ or $[150m, 240m]$), the accuracy is not highly




FIG. 11: Error of amplitude extrapolation to infinity for extrapolation with order n , cf. Eq. (20). Plotted are relative amplitude differences between extrapolation with orders n and $n+1$. The inset is an enlargement for $t - r^* \gtrsim 1000m$.

variable, though we find that noise is increased when using data from such a smaller range of extraction radii.

To estimate the errors generated by not extrapolating waveforms to infinity at all, Fig. 12 contains also the phase difference between wave extraction at two finite radii ($90m$ and $240m$) and our preferred extrapolated phase at infinity. The dotted lines show such phase differences when only a time shift by the tortoise-coordinate radius of the extraction sphere is applied. The errors are dramatic, tenths of radians or more, even very late in the run. When matching to post-Newtonian waveforms, we are free to add an overall time and phase shift (cf. Section IV B). Therefore, the dashed lines in Fig. 12 show phase differences with the same unextrapolated waveforms as shown by the dotted lines, except that a phase and time shift has been applied so that the ϕ and $\dot{\phi}$ agree with those of the extrapolated waveform late in the run (where $m\omega = 0.1$), where the wavelengths are shortest and wave extraction is expected to work best. Even with such an adjustment, the gravitational wave phase extracted at $r = 90m$ differs by about 0.1 radian at $t \sim 1000m$ before coalescence, with this difference growing to 0.3 radians at the start of our simulation.

Figure 13 makes the same comparison for the gravitational wave amplitude. Wave extraction at $r = 90m$ results in relative amplitude errors of up to 8 per cent, and of about 2 per cent even in the last 1000m of our simulation. We also point out that the errors due to finite extraction radius decay approximately as the inverse of the extraction radius: For waves extracted at $r = 240m$ the errors are smaller than for waves extracted at $r = 90m$ by about a factor of three, as can be seen in Figs. 12 and 13; for wave extraction at $r = 45m$, the errors would be approximately twice as large as the $r = 90m$ case.




FIG. 12: Effect of wave extraction radii on extrapolated phase. Each curve represents the difference from our preferred wave extrapolation using $r \in [75m, 240m]$. The three solid curves represent extrapolation from different intervals of extraction radii. The curves labeled “240m” and “90m” represent differences from waves extracted at these two radii, without any extrapolation, for two cases: time and phase shifted so that ϕ and $\dot{\phi}$ match at $m\omega = 0.1$ (dashed), and without these shifts (dotted).




FIG. 13: Effect of choice of wave extraction radii on extrapolated amplitude. Each curve represents the (relative) amplitude difference to our preferred wave extrapolation using $r \in [75m, 240m]$. The three solid curves represent extrapolation from different intervals of extraction radii. The curves labeled “240m” and “90m” represent differences from waves extracted at these two radii, without any extrapolation, for two cases: time and phase shifted so that ϕ and $\dot{\phi}$ match at $m\omega = 0.1$ (dashed), and without these shifts (dotted).

The errors introduced by using a finite extraction radius are significantly larger than our truncation error (even at extraction radius $240m$). Therefore extrapolation to infinity is essential to realize the full accuracy of our simulations.

III. GENERATION OF POST-NEWTONIAN WAVEFORMS

It is not our intention to review all of post-Newtonian (PN) theory, but to summarize the important points that go into the construction of the post-Newtonian waveforms that we will compare to our numerical simulation. For a complete review of post-Newtonian methods applied to inspiralling compact binaries, see the review article by Blanchet[99].

The post-Newtonian approximation is a slow-motion, weak-field approximation to general relativity with an expansion parameter $\epsilon \sim (v/c)^2 \sim (Gm/rc^2)$. For a binary system of two point masses m_1 and m_2 , v is the magnitude of the relative velocity, m is the total mass, and r is the separation. In order to produce a post-Newtonian waveform, it is necessary to solve both the post-Newtonian equations of motion describing the binary, and the post-Newtonian equations describing the generation of gravitational waves.

Solving the equations of motion yields explicit expressions for the accelerations of each body in terms of the positions and velocities of the two bodies[100–110]. The two-body equations of motion can then be reduced to relative equations of motion in the center-of-mass frame in terms of the relative position and velocity[111]. The relative acceleration is currently known through 3.5PN order, where 0PN order for the equations of motion corresponds to Newtonian gravity. The effects of radiation reaction (due to the emission of gravitational waves) enters the relative acceleration starting at 2.5PN order. The relativistic corrections to the relative acceleration at 1PN, 2PN and 3PN order (ignoring the radiation reaction terms at 2.5PN and 3.5PN order) admit a conserved center of mass binding energy through 3PN order[112]. There is no 2.5PN or 3.5PN order contribution to the energy.

Solving the post-Newtonian wave generation problem yields expressions for the gravitational waveform h_{ij} and gravitational wave flux \mathcal{L} in terms of radiative multipole moments[113]. These radiative multipole moments are in turn related to the source multipole moments, which can be given in terms of the relative position and relative velocity of the binary[114]. For the gravitational wave generation problem, PN orders are named with respect to the leading order waveform and flux, which are given by the quadrupole formalism. Thus, for example, 1.5PN order in the wave generation problem represents terms of order $(v/c)^3$ beyond quadrupole. Higher order effects enter both through post-Newtonian corrections to the mass quadrupole, as well as effects due to higher multipole mo-

ments. Starting at 1.5PN order the radiative multipole moments include non-linear and non-instantaneous (i.e. depend upon the past history of the binary) interactions among the source multipole moments (e.g. gravitational wave tails)[114–117].

It was recognized early that simply plugging in the orbital evolution predicted by the equations of motion into the expressions for the waveform would not generate templates accurate enough for matched filtering in detecting gravitational waves[118]. This is because radiation reaction enters the equations of motion only at the 2.5PN order; hence computing a waveform to k PN order beyond the quadrupole formalism would require $2.5 + k$ PN orders in the equations of motion. In order to obtain as accurate a post-Newtonian waveform as possible it is thus necessary to introduce the assumption of an adiabatic inspiral of a quasi-circular orbit, as well as the assumption of energy balance between the orbital binding energy and the energy emitted by the gravitational waves.

A. Adiabatic inspiral of quasi-circular orbits

The emission of gravitational radiation causes the orbits of an isolated binary system to circularize [48]. Thus it is a reasonable assumption to model the orbital evolution of the binary as a slow adiabatic inspiral of a quasi-circular orbit. With this assumption, post-Newtonian expressions for the orbital energy E and gravitational energy flux \mathcal{L} are currently known through 3.5PN order [119–123]. These expressions can be given in terms of a parameter related to either the harmonic coordinate separation r , or to the orbital frequency Ω . We choose to use the expressions given in terms of a frequency-related parameter

$$x \equiv \left(\frac{Gm\Omega}{c^3} \right)^{2/3} \quad (23)$$

rather than a coordinate-related parameter, because the coordinate relationship between the numerical simulation and the harmonic coordinates used in post-Newtonian approximations is unknown. The orbital energy for an equal mass system is given by[99]

$$E = -\frac{mc^2}{8}x \left[1 - \frac{37}{48}x - \frac{1069}{384}x^2 + \left(\frac{1427365}{331776} - \frac{205}{384}\pi^2 \right) x^3 \right], \quad (24)$$

and the gravitational wave flux for an equal mass system is given by [99]

$$\mathcal{L} = \frac{2c^5}{5G}x^5 \left\{ 1 - \frac{373}{84}x + 4\pi x^{3/2} - \frac{59}{567}x^2 - \frac{767}{42}\pi x^{5/2} + \left[\frac{18608019757}{209563200} + \frac{355}{64}\pi^2 - \frac{1712}{105}\gamma - \frac{856}{105}\ln(16x) \right] x^3 + \frac{16655}{6048}\pi x^{7/2} \right\}, \quad (25)$$

where $\gamma = 0.577216\dots$ is Euler's constant.

B. Polarization Waveforms

The gravitational polarization waveforms for a quasi-circular orbit in the $x - y$ plane, as measured by an observer at spherical coordinates $(R, \hat{\theta}, \hat{\phi})$, are given by

$$h_+ = \frac{2G\mu}{c^2R}x \left\{ -(1 + \cos \hat{\theta}) \cos 2(\Phi - \hat{\phi}) + \dots \right\} \quad (26)$$

$$h_\times = \frac{2G\mu}{c^2R}x \left\{ -2 \cos \hat{\theta} \sin 2(\Phi - \hat{\phi}) + \dots \right\}, \quad (27)$$

where Φ is the orbital phase (measured from the x-axis) and $\mu = m_1 m_2 / m$ is the reduced mass. The polarization waveforms are currently known through 2.5PN order[124, 125].

1. Optimally oriented observer

For an equal-mass binary the polarization waveforms along the z-axis (i.e. the optimally oriented observer along the normal to the orbital plane) are given by [124, 125]

$$h_+^{(z)} = \frac{Gm}{2c^2R}x \left(\cos 2\Phi \left\{ -2 + \frac{17}{4}x - 4\pi x^{3/2} + \frac{15917}{2880}x^2 + 9\pi x^{5/2} \right\} \right. \\ \left. + \sin 2\Phi \left\{ -12 \ln \left(\frac{x}{x_0} \right) x^{3/2} + \left[\frac{59}{5} + 27 \ln \left(\frac{x}{x_0} \right) \right] x^{5/2} \right\} \right) \quad (28)$$

$$h_\times^{(z)} = \frac{Gm}{2c^2R}x \left(\sin 2\Phi \left\{ -2 + \frac{17}{4}x - 4\pi x^{3/2} + \frac{15917}{2880}x^2 + 9\pi x^{5/2} \right\} \right. \\ \left. + \cos 2\Phi \left\{ 12 \ln \left(\frac{x}{x_0} \right) x^{3/2} - \left[\frac{59}{5} + 27 \ln \left(\frac{x}{x_0} \right) \right] x^{5/2} \right\} \right), \quad (29)$$

where

$$\ln x_0 \equiv \frac{11}{18} - \frac{2}{3}\gamma + \frac{2}{3} \ln \left(\frac{Gm}{4bc^3} \right) \quad (30)$$

is a constant frequency scale that depends upon the constant time scale b entering the gravitational wave tail contribution to the polarization waveforms [126, 127]. The freely-specifiable constant b corresponds to a choice of the origin of radiative time T with respect to harmonic time t , and enters the relation between the retarded time $T_R = T - R/c$ in radiative coordinates (the coordinates in which the waveform is given) and the retarded time $t - r/c$ in harmonic coordinates (the coordinates in which the equations of motion are given) [126, 127]:

$$T_R = t - \frac{r}{c} - \frac{2GM_{\text{ADM}}}{c^3} \ln \left(\frac{r}{bc} \right). \quad (31)$$

Here M_{ADM} is the ADM mass (mass monopole) of the binary system.

2. The (2,2) mode

When comparing a post-Newtonian waveform with data from a physical gravitational wave detector, it is necessary to compare waves emitted in a certain direction $(\hat{\theta}, \hat{\phi})$ with respect to the source. However, comparing waveforms between PN and numerical simulations can be done in all directions simultaneously by decomposing the waveforms in terms of spherical harmonics and then comparing different spherical harmonic modes. Since the power in each spherical harmonic mode decreases rapidly with spherical harmonic index, with the (2, 2) mode dominating (for an equal-mass non-spinning binary), it is possible to do a very accurate comparison that is valid for all angles by using only a few modes. In addition, as pointed out by Kidder [60], the dominant (2,2) mode can be computed to 3PN order. For an equal-mass binary, the (2,2) mode is

$$h_{(2,2)} = -2\sqrt{\frac{\pi}{5}} \frac{Gm}{c^2R} e^{-2i\Phi} x \left\{ 1 - \frac{373}{168}x + \left[2\pi + 6i \ln \left(\frac{x}{x_0} \right) \right] x^{3/2} - \frac{62653}{24192}x^2 - \left[\frac{197}{42}\pi + \frac{197i}{14} \ln \left(\frac{x}{x_0} \right) + 6i \right] x^{5/2} \right. \\ \left. + \left[\frac{43876092677}{1117670400} + \frac{99}{128}\pi^2 - \frac{428}{105} \ln x - \frac{856}{105}\gamma - \frac{1712}{105} \ln 2 - 18 \left[\ln \left(\frac{x}{x_0} \right) \right]^2 + \frac{428}{105}i\pi + 12i\pi \ln \left(\frac{x}{x_0} \right) \right] x^3 \right\}. \quad (32)$$

Since the (2,2) mode of the numerical waveforms is less noisy than the waveform measured along the z-axis, and since we have access to the 3PN amplitude correction of the (2,2) mode, we will use the (2,2) waveforms rather than the z-axis waveforms for our comparisons between NR and PN in Sec. VI. We have verified (for all comparisons using post-Newtonian waveforms of ≤ 2.5 PN order in amplitude) that our results do not change significantly when we use z-axis waveforms instead of (2,2) waveforms.

C. Absorbing amplitude terms into a redefinition of the phase

The logarithms of the orbital frequency parameter x (as well as the constant frequency scale x_0) that appear in the amplitude expressions (28), (29), and (32) can be absorbed into a redefinition of the phase by introducing an auxiliary phase variable $\Psi = \Phi + \delta$. Noting that the $\ln x$ terms first enter at 1.5 PN order, it is straightforward to show that choosing [60, 124, 128]

$$\delta = -3 \frac{M_{\text{ADM}}}{m} x^{3/2} \ln \left(\frac{x}{x_0} \right), \quad (33)$$

where $M_{\text{ADM}}/m = 1 - x/8 + O(x^2)$ for an equal mass system, will eliminate the $\ln x$ terms from both the (2,2) mode as well as for the polarization waveforms. This follows from

$$\begin{aligned} h_{(2,2)} &= Ae^{-2i\Psi} \\ &= Ae^{-2i\Phi} e^{-2i\delta} \\ &= Ae^{-2i\Phi}(1 - 2i\delta - 2\delta^2 + O(x^{9/2})), \end{aligned}$$

and similarly for the polarization waveforms. Furthermore, since the orbital phase as a function of frequency goes as $x^{-5/2}$ at leading order (see Eq. (40) below), the $\ln x$ terms, which were 1.5PN, 2.5PN, and 3PN order in the original amplitude expressions, now appear as phase corrections at relative order 4PN, 5PN, and 5.5PN. As these terms are beyond the order to which the orbital phase evolution is known (3.5PN order), it can be argued that these terms can be ignored. Note that the choices of x_0 in Eq. (30) and δ in Eq. (33) are not unique; they were made to gather all logarithmic terms into one term, as well as to simplify the waveform [128].

D. Energy balance

The second assumption that goes into making as accurate a post-Newtonian waveform as possible is that of energy balance. It is assumed that the energy carried away by the emission of gravitational waves is balanced by the change in the orbital binding energy of the binary,

$$\frac{dE}{dt} = -\mathcal{L}. \quad (34)$$

While this is extremely plausible, it has only been confirmed through 1.5 PN order [129].

Given the above expressions for the energy, flux, and waveform amplitude, there is still a set of choices that must be made in order to produce a post-Newtonian waveform that can be compared to our numerical waveform. These include

1. The PN order through which terms in the orbital energy and luminosity are retained.
2. The procedure by which the energy balance equation is used to obtain $x(t)$ and $\Phi(t)$.
3. The PN order through which terms in the waveform amplitude are kept.
4. The treatment of the $\ln x$ terms. These terms can be included in the amplitude or included in the orbital phase via the auxiliary phase $\Psi \equiv \Phi + \delta$. If the latter is chosen, these terms can be retained or ignored; ignoring them can be justified because they occur at higher order than all known terms in the orbital phase.

Unless explicitly stated otherwise, we will choose to retain terms through 3.5PN order in the orbital energy and luminosity, to retain terms through 2.5PN order in the waveform amplitude, and will ignore the $\ln x$ terms in the amplitude by absorbing them into the phase and dropping them because of their high PN order. In the next section we describe several choices for obtaining $x(t)$ and $\Phi(t)$ from the energy balance equation.

E. Taylor approximants: Computing $\Phi(t)$

In this section we describe how to obtain the orbital phase as a function of time, $\Phi(t)$, using the energy balance equation (34). Different methods of doing this exist; here we follow the naming convention of [52]. These methods, and variations of them, are called Taylor approximants, and all formally agree to a given PN order but differ in how higher-order terms are truncated. We discuss four time-domain approximants here, but more can be defined.

1. TaylorT1

The TaylorT1 approximant is obtained by numerically integrating the ODEs

$$\frac{dx}{dt} = -\frac{\mathcal{L}}{(dE/dx)} \quad (35)$$

$$\frac{d\Phi}{dt} = \frac{c^3}{Gm} x^{3/2}, \quad (36)$$

to produce $\Phi(t)$. The fraction on the right side of Eq (35) is retained as a ratio of post-Newtonian expansions, and is not expanded further before numerical integration. This is the approximant used in the NR-PN comparisons in [33, 39].

2. TaylorT2

The TaylorT2 approximant is obtained by starting with the parametric solution of the energy balance equation:

$$t(x) = t_0 + \int_x^{x_0} dx \frac{(dE/dx)}{\mathcal{L}} \quad (37)$$

$$\Phi(x) = \Phi_0 + \int_x^{x_0} dx \frac{x^{3/2} c^3}{Gm} \frac{(dE/dx)}{\mathcal{L}}. \quad (38)$$

The integrand of each expression is re-expanded as a single post-Newtonian expansion in x and truncated at the appropriate PN-order; these integrals are then evaluated

analytically to obtain for an equal-mass binary [52, 53]:

$$\begin{aligned}
t &= t_0 - \frac{5Gm}{64c^3} x^{-4} \left\{ 1 + \frac{487}{126} x - \frac{32}{5} \pi x^{3/2} + \frac{2349439}{254016} x^2 \right. \\
&\quad \left. - \frac{1864}{63} \pi x^{5/2} + \left[-\frac{999777207379}{5867769600} + \frac{1597}{48} \pi^2 \right. \right. \\
&\quad \left. \left. + \frac{6848}{105} \gamma + \frac{3424}{105} \ln(16x) \right] x^3 - \frac{571496}{3969} \pi x^{7/2} \right\} \quad (39) \\
\Phi &= \Phi_0 - \frac{1}{8} x^{-5/2} \left\{ 1 + \frac{2435}{504} x - 10\pi x^{3/2} + \frac{11747195}{508032} x^2 \right. \\
&\quad \left. + \frac{1165}{42} \pi x^{5/2} \ln x + \left[\frac{1573812724819}{4694215680} - \frac{7985}{192} \pi^2 \right. \right. \\
&\quad \left. \left. - \frac{1712}{21} \gamma - \frac{856}{21} \ln(16x) \right] x^3 + \frac{357185}{7938} \pi x^{7/2} \right\}. \quad (40)
\end{aligned}$$

3. TaylorT3

The TaylorT3 approximant is closely related to TaylorT2. It is obtained by introducing the dimensionless time variable

$$\tau \equiv \frac{\nu c^3}{5Gm} (t_0 - t), \quad (41)$$

where $\tau^{-1/4} = O(\epsilon)$. The TaylorT2 expression $t(x)$ is inverted to obtain $x(\tau)$, and truncated at the desired PN order. Then $x(\tau)$ is integrated to obtain

$$\Phi(\tau) = \Phi_0 - \int_{\tau_0}^{\tau} d\tau \frac{5x^{3/2}}{\nu}. \quad (42)$$

This procedure yields for an equal-mass binary [99]:

$$\begin{aligned}
x &= \frac{1}{4} \tau^{-1/4} \left\{ 1 + \frac{487}{2016} \tau^{-1/4} - \frac{1}{5} \pi \tau^{-3/8} \right. \\
&\quad + \frac{1875101}{16257024} \tau^{-1/2} - \frac{1391}{6720} \pi \tau^{-5/8} \\
&\quad + \left[-\frac{999777207379}{1502149017600} + \frac{1597}{12288} \pi^2 + \frac{107}{420} \gamma \right. \\
&\quad \left. - \frac{107}{3360} \ln\left(\frac{\tau}{256}\right) \right] \tau^{-3/4} - \frac{88451}{282240} \pi \tau^{-7/8} \left. \right\} \quad (43) \\
\Phi &= \Phi_0 - 4\tau^{5/8} \left\{ 1 + \frac{2435}{4032} \tau^{-1/4} - \frac{3}{4} \pi \tau^{-3/8} \right. \\
&\quad + \frac{1760225}{1806336} \tau^{-1/2} - \frac{1165}{5376} \pi \tau^{-5/8} \ln \tau \\
&\quad + \left[\frac{24523613019127}{3605157642240} - \frac{42997}{40960} \pi^2 - \frac{107}{56} \gamma \right. \\
&\quad \left. + \frac{107}{448} \ln\left(\frac{\tau}{256}\right) \right] \tau^{-3/4} + \frac{28325105}{21676032} \pi \tau^{-7/8} \left. \right\} \quad (44)
\end{aligned}$$

This is the post-Newtonian approximant used in visual comparisons by [32] at 3PN order in phase.

4. TaylorT4

In addition to simply numerically integrating the flux-energy equation (35), as is done for TaylorT1, one may instead re-expand the right side of (35) as a single series and truncate at the appropriate PN order before doing the integration. The phase evolution $\Phi(t)$ can thus be obtained by numerically integrating the ODEs

$$\begin{aligned}
\frac{dx}{dt} &= \frac{16c^3}{5Gm} x^5 \left\{ 1 - \frac{487}{168} x + 4\pi x^{3/2} + \frac{274229}{72576} x^2 \right. \\
&\quad - \frac{254}{21} \pi x^{5/2} + \left[\frac{178384023737}{3353011200} + \frac{1475}{192} \pi^2 - \frac{1712}{105} \gamma \right. \\
&\quad \left. \left. - \frac{856}{105} \ln(16x) \right] x^3 + \frac{3310}{189} \pi x^{7/2} \right\} \quad (45)
\end{aligned}$$

$$\frac{d\Phi}{dt} = \frac{x^{3/2} c^3}{Gm}. \quad (46)$$

This approximant was not considered in [52], however for consistency with their notation, we call it TaylorT4. TaylorT4 is the primary approximant used in the NR-PN comparisons in [37, 38], and one of the several approximants considered in the NR-PN comparisons in [33]. Ref.[32] pointed out that TaylorT4 at 3.5PN order in phase is close to TaylorT3 at 3PN order in phase, and therefore should give similar agreement with numerical results.

IV. PN-NR COMPARISON PROCEDURE

A. What to compare?

There are many ways to compare numerical relativity and post-Newtonian results. For example, the post-Newtonian orbital phase $\Phi(t)$ could be compared with the coordinate phase of the black hole trajectories. However, this and many other comparisons are difficult to make in a coordinate-independent manner without expending significant effort to understand the relationship between the gauge choices used in post-Newtonian theory and in the NR simulations. Therefore, in order to obtain the most meaningful comparison possible, we attempt to minimize gauge effects by comparing gravitational waveforms as seen by an observer at infinity. The waveform quantity most easily obtained from the numerical relativity code is the Newman-Penrose quantity Ψ_4 , and we will predominantly compare its $l=2, m=2$ component [cf. Eq. (13)], split into phase ϕ and amplitude A according to Eq. (14) and extrapolated to infinite extraction radius.

The post-Newtonian formulae in Section III yield the metric perturbation components h_+ and h_\times , which—for a gravitational wave at infinity—are related to Ψ_4 by

$$\Psi_4 = \frac{\partial^2}{\partial t^2} (h_+ - ih_\times). \quad (47)$$

We numerically differentiate the post-Newtonian expressions for h_+ and h_\times twice before computing amplitude

and phase using Eq. (14). Note that $\phi(t)$ will differ slightly from the phase computed from the metric perturbation directly, as $\tan^{-1}(h_{\times}/h_{+})$, because both the amplitude and phase of the metric perturbation are time dependent. For the same reason, $\phi(t)$ is not precisely equal to twice the orbital phase.

As in Ref. [39], we compare Ψ_4 rather than $h_{+,\times}$ to avoid difficulties arising with fixing the integration constants when integrating the numerically obtained Ψ_4 (see [36] for more details). Both Ψ_4 and $h_{+,\times}$ contain the same information, so differences between both procedures should be minimal.

B. Matching procedure

Each of the post-Newtonian waveforms has an arbitrary time offset t_0 and an arbitrary phase offset ϕ_0 . These constants can be thought of as representing the absolute time of merger and the orientation of the binary at merger, and we are free to adjust them in order to match NR and PN waveforms. Following [37, 39], we choose these constants by demanding that the PN and NR gravitational wave phase and gravitational wave frequency agree at some fiducial frequency ω_m . Specifically, we proceed as follows: We start with a NR waveform $\Psi_4^{\text{NR}}(t)$ and an unshifted PN waveform $\Psi_4^{\text{PN}'}(t)$ that has an arbitrary time and phase shift. After selecting the matching frequency ω_m , we can find (to essentially unlimited accuracy) the time t_c such that the derivative of the PN phase satisfies $\dot{\phi}_{\text{PN}'}(t_c) = \omega_m$, where $\phi_{\text{PN}'}(t)$ is the phase associated with $\Psi_4^{\text{PN}'}(t)$. Similarly, we find the time t_m such that $\dot{\phi}_{\text{NR}}(t_m) = \omega_m$. The time t_m cannot be found to unlimited accuracy, and the uncertainty in t_m is due mainly to residual eccentricity of the NR waveform, as discussed in Section V E. Once we have t_m and t_c , we leave the NR waveform untouched, but we construct a new, shifted, PN waveform

$$\Psi_4^{\text{PN}}(t) = \Psi_4^{\text{PN}'}(t + t_c - t_m)e^{i(\phi_{\text{PN}'}(t_c) - \phi_{\text{NR}}(t_m))}. \quad (48)$$

The phase of this new PN waveform is therefore

$$\phi_{\text{PN}}(t) = \phi_{\text{PN}'}(t + t_c - t_m) - \phi_{\text{PN}'}(t_c) + \phi_{\text{NR}}(t_m), \quad (49)$$

which satisfies $\phi_{\text{PN}}(t_m) = \phi_{\text{NR}}(t_m)$ and $\dot{\phi}_{\text{PN}}(t_m) = \omega_m$ as desired. All our comparisons are then made using the new shifted waveform $\Psi_4^{\text{PN}}(t)$ rather than the unshifted waveform $\Psi_4^{\text{PN}'}(t)$.

C. Choice of Masses

The post-Newtonian expressions as written in Section III involve the total mass m , which corresponds to the the sum of the bare masses of the point particles in post-Newtonian theory. When comparing PN to NR, the question then arises as to which of the many definitions of

the mass of a numerically-generated binary black hole solution should correspond to the post-Newtonian parameter m . For non-spinning black holes at very large separation, m reduces to the sum of the irreducible masses of the two holes. Neglecting tidal heating, the irreducible masses should be conserved during the inspiral, so that we identify m with the sum of the irreducible masses of the initial data 30c. As discussed in Sec. V the black hole spins are sufficiently small so that there is no discernible difference between irreducible mass of the black holes and the Christoudoulou mass, Eq. (3). Of course, the latter would be more appropriate for spinning black holes.

V. ESTIMATION OF UNCERTAINTIES

To make precise statements about agreement or disagreement between numerical and post-Newtonian waveforms, it is essential to know the size of the uncertainties in this comparison. When discussing these uncertainties, we must strive to include all effects that may cause our numerical waveform to differ from the post-Newtonian waveforms we compare to. For instance, in addition to considering effects such as numerical truncation error, we also account for the fact that NR and PN waveforms correspond to slightly different physical scenarios: The PN waveforms have identically zero spin and eccentricity, whereas the numerical simulations have some small residual spin and eccentricity. Table III lists all effects we have considered; we discuss these in detail below starting in Sec. V A). All uncertainties are quoted in terms of phase and amplitude differences, and apply to waveform comparisons via matching at a fixed ω_m according to the procedure in Sec. IV B.

Most of the effects responsible for our uncertainties are time dependent, so that it is difficult to arrive at a single number describing each effect. For simplicity, the error bounds in Table III ignore the junk-radiation noise that occurs in the numerical waveform for $t - r^* \lesssim 1000m$. The extent to which this noise affects the PN-NR comparisons presented below in Sections VIA and VIB will be evident from the noise in the graphs in these sections. Note that all four matching frequencies ω_m occur after the noise disappears at $t - r^* \sim 1000m$. Furthermore, the post-Newtonian waveforms end at different times depending on the PN order and on which particular post-Newtonian approximant is used. Therefore, in order to produce a single number for each effect listed in Table III, we consider only the part of the waveform prior to some cutoff time, which we choose to be the time at which the numerical waveform reaches gravitational wave frequency $m\omega = 0.1$.

TABLE III: Summary of uncertainties in the comparison between numerical relativity and post-Newtonian expansions. Quoted error estimates ignore the junk-radiation noise at $t \lesssim 1000m$ and apply to times before the numerical waveform reaches gravitational wave frequency $m\omega = 0.1$. Uncertainties apply to waveform comparisons via matching at a fixed ω_m according to the procedure in Sec. IV B, and represent the maximum values for all four different matching frequencies ω_m that we consider, unless noted otherwise.

Effect	$\delta\phi$ (radians)	$\delta A/A$
Numerical truncation error	0.003	0.001
Finite outer boundary	0.005	0.002
Extrapolation $r \rightarrow \infty$	0.005	0.002
GW extraction at $r_{\text{areal}}=\text{const?}$	0.002	10^{-4}
Drift of mass m	0.002	10^{-4}
Coordinate time = proper time?	0.002	10^{-4}
Lapse spherically symmetric?	0.01	4×10^{-4}
residual eccentricity	0.02^a	0.004
residual spins	0.03	0.001
root-mean-square sum	0.04^a	0.005

^aFor the case of matching at $m\omega_m = 0.04$, the phase uncertainty due to residual eccentricity increases to 0.05 radians, thus increasing the root-mean-square sum to 0.06 radians.

A. Errors in numerical approximations

The first three error sources listed in Table III have already been discussed in detail in Section II. We estimate numerical truncation error using the difference between the two highest resolution runs after the waveforms have been shifted to agree at some matching frequency ω_m . For $m\omega_m = 0.1$ this difference is shown as the curves labeled '30c-1/N5' in the lower panels of Figs. 7 and 8, and corresponds to a phase difference of 0.003 radians and a relative amplitude difference of 0.001. For other values of ω_m the differences are similar. The effect of the outer boundary is estimated by the difference between the runs 30c-1/N6 and 30c-2/N6, which for $m\omega_m = 0.1$ is shown as the curves labeled '30c-2/N6' in the lower panels of Figs. 7 and 8, and amount to phase differences of 0.005 radians and relative amplitude differences of 0.002. Errors associated with extrapolation to infinity have been discussed in detail in Figs. 10 and 12. Specifically, Fig. 10 shows that increasing the extrapolation order between 3 and 4 changes the extrapolated phase by less than 0.005 radians, and Fig. 12 confirms that the extrapolated result is robust under changes of extraction radii.

B. Constancy of extraction radii

If the physical locations of the coordinate-stationary extraction radii happen to change during the evolution, then the extracted gravitational waves will accrue a timing error equal to the light-travel time between the original location and the final location. From Fig. 9, we see

that the drift in areal radius is less than $0.02m$, resulting in a time uncertainty of $\delta t = 0.02m$. This time uncertainty translates into a phase uncertainty via

$$\delta\phi = m\omega \times (\delta t/m) \quad (50)$$

which yields $\delta\phi \approx 0.002$, when $m\omega = 0.1$ (the value at the end of the PN comparison) was used.

To estimate the effect of this time uncertainty on the amplitude, we first note that to lowest order in the post-Newtonian parameter x (defined in Eq. (23)), the wave amplitude of Ψ_4 scales like x^4 . Also, from Eq. (45), we have $dx/dt = 16/(5m)x^5$. Therefore,

$$\frac{\delta A}{A} \sim \frac{d \ln A}{dx} \frac{dx}{dt} \delta t \sim \frac{64}{5} (m\omega/2)^{8/3} \frac{\delta t}{m}, \quad (51)$$

where we have used the fact that the gravitational wave frequency ω is approximately twice the orbital frequency. For a time uncertainty $\delta t = 0.02m$, Eq. (51) gives $\delta A/A \approx 10^{-4}$ for $m\omega = 0.1$.

C. Constancy of mass

Our comparisons with post-Newtonian formulae assume a constant post-Newtonian mass parameter m , which we set equal to the total irreducible mass of the black holes in the numerical simulation. If the total mass of the numerical simulation is not constant, this will lead to errors in the comparison. For example, changes in t/m caused by a changing mass will lead to phase differences. Figure 4 demonstrates that the irreducible mass is conserved to a fractional accuracy of about $\delta m/m \approx 5 \times 10^{-6}$. For $t/m = 4000$, this gives an uncertainty in the overall time interval of $\delta(t/m) = (t/m) \times (\delta m/m) \approx 0.02$. This time uncertainty translates into a phase uncertainty of $\delta\phi \approx 0.002$, using Eq. (50) for $m\omega = 0.1$. Note that the effect of the black-hole spins on the mass is negligible relative to the numerical drift of 5×10^{-6} . This is because the spins of the holes are bounded by $S/M_{\text{irr}} < 2 \times 10^{-4}$ and the spin enters quadratically into the Christoudoulou formula (3). The error in the gravitational wave amplitude caused by time uncertainties due to varying mass is $\delta A/A \approx 10^{-4}$ using Eq. (51) for $m\omega = 0.1$. An error in the mass will affect the amplitude not only via a time offset, but also because the amplitude is proportional to $(\omega m/2)^{8/3}$ (to lowest PN order). However, this additional error is very small, $\delta A/A \approx (8/3)\delta m/m \approx 10^{-5}$.

D. Time coordinate ambiguity

We now turn to two possible sources of error that have not yet been discussed, both of which are related to ambiguity in the time coordinate. The basic issue is that the time variable t in post-Newtonian expansions corresponds to proper time in the asymptotically flat region, but the time t in numerical simulations is coordinate time. These two quantities agree only if the lapse




FIG. 14: Asymptotic behavior of the lapse at large radii for times $t/m = 0, 1900, 3800$. The top figure displays the angular average of the lapse as a function of radius at $t = 0, 1900m, 7800m$. The bottom figure shows the dominant higher multipole moments of the lapse. Both horizontal axes are spaced in $1/r$.

function N approaches unity at large distances. To verify this, we decompose N in spherical harmonics centered on the center of mass of the system,

$$N(r, \theta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l N_{lm}(r) Y_{lm}(\theta, \varphi). \quad (52)$$

The angular average of the lapse function, $\bar{N}(r) \equiv \sqrt{4\pi} N_{00}$ should then approach unity for $r \rightarrow \infty$, and all other modes $N_{lm}(r)$ should decay to zero. The top panel of Fig. 14 plots $\bar{N}(r) - 1$ vs. m/r for three different evolution times. Fitting $\bar{N}(r) - 1$ for $r > 100m$ to a polynomial in m/r gives a y-intercept of $< 5 \times 10^{-6}$ for all three times, and for polynomial orders of two through five. Therefore, the coordinate time of the evolution agrees with proper time at infinity to better than $\delta t/m = t/m \times 5 \times 10^{-6} \approx 0.02$, which induces a phase error of at most $\delta\phi \approx 0.002$ and an amplitude error of $\delta A/A \approx 10^{-4}$.

The second source of error related to the lapse is shown in the lower panel of Fig. 14, which presents the three dominant higher order moments $N_{lm}(r)$. All these modes decay to zero as $r \rightarrow \infty$, except, perhaps, the real part of the N_{22} mode at $t/m = 3800$. This mode seems to approach a value of about 5×10^{-5} . At $t = 1900m$, this mode still decays nicely to zero, hence the maximum time uncertainty introduced by this effect at late times is $\delta t = 1900m \times 5 \times 10^{-5} \approx 0.1m$, resulting in a potential phase uncertainty of $\delta\phi \approx 0.01$ and a potential amplitude uncertainty of $\delta A/A \approx 4 \times 10^{-4}$.

E. Eccentricity

We estimated the eccentricity during the numerical simulation with several of the methods described in [32, 36, 59], and have found consistently $e \lesssim 6 \times 10^{-5}$. This eccentricity can affect our comparison to a post-Newtonian waveform of a quasi-circular (i.e. zero eccentricity) inspiral in three ways.

1. Change in rate of inspiral

The first effect arises because an eccentric binary has a different inspiral rate than a non-eccentric binary; physically, this can be understood by noting that the gravitational flux and orbital energy depend upon the eccentricity, and therefore modify the rate at which the orbital frequency evolves assuming energy balance. Reference [130] has derived the first-order correction in the phase of the gravitational wave due to this effect. Converting their result to our notation and restricting to the equal mass case yields

$$\frac{1}{(dx/dt)} = \frac{5Gm}{16c^3 x^5} \left(1 - \frac{157}{24} e_i^2 \left(\frac{x_i}{x} \right)^{19/6} \right), \quad (53)$$

where e_i is the initial eccentricity and x_i is the initial value of the orbital frequency parameter. Substituting this into Eq. (38) yields

$$\Phi = \Phi_0 - \frac{1}{8} x^{-5/2} + \frac{785}{2176} e_i^2 x_i^{19/6} x^{-17/3}. \quad (54)$$

Using $e_i = 6 \times 10^{-5}$ and integrating over the frequency range from the start of our simulation to the matching frequency of $m\omega = 0.1$ yields a phase shift of $\sim -2 \times 10^{-6}$, which is dwarfed by many other error sources, such as the uncertainty in the numerical mass m , cf. Sec. V C.

2. Uncertainty in matching time

The second way in which eccentricity affects our comparison is by introducing errors in our procedure for matching the PN and NR waveforms. Recall that our matching procedure involves determining the time t_m at which the gravitational wave frequency ω takes a certain value $m\omega_m$; eccentricity modulates the instantaneous gravitational wave frequency $\omega(t)$ via

$$\omega(t) = \bar{\omega}(t) [1 + 2e \cos(\Omega_r t)], \quad (55)$$

where $\bar{\omega}(t)$ represents the averaged “non-eccentric” evolution of the gravitational wave frequency, and Ω_r is the frequency of radial oscillations, which is approximately the orbital frequency. We see that ω can differ from $\bar{\omega}$ by as much as $2e\bar{\omega} \approx 2e\omega$. This could induce an error in determination of t_m by as much as

$$|\delta t_m| = \frac{|\delta\omega|}{\dot{\omega}} \approx \frac{2e\omega}{\dot{\omega}} \quad (56)$$

We can simplify this expression by using Eq. (45) to lowest order, and by noting that the gravitational wave frequency is approximately twice the orbital frequency. We find

$$|\delta t_m| \leq e \frac{5m}{12} \left(\frac{m\omega}{2}\right)^{-8/3}. \quad (57)$$

This uncertainty is largest at *small* frequencies, because the frequency changes much more slowly. For $m\omega = 0.04$, we find $|\delta t_m| \lesssim 0.9m$, and for $m\omega = 0.1$, we find $|\delta t_m| \lesssim 0.1m$.

To determine how uncertainties in t_m translate into phase differences, recall that in the matching procedure described in Section IV B, t_m enters into the phase of the shifted PN waveform according to Eq. (49). Therefore the phase difference that we compute between the PN and NR waveforms is

$$\begin{aligned} \Delta\phi(t) &= \phi_{\text{PN}}(t) - \phi_{\text{NR}}(t) \\ &= \dot{\phi}_{\text{PN}'}(t+t_c-t_m) - \dot{\phi}_{\text{NR}}(t) + \phi_{\text{NR}}(t_m) - \dot{\phi}_{\text{PN}'}(t_c). \end{aligned} \quad (58)$$

Then the error in $\Delta\phi$ is found by Taylor expanding Eq. (58):

$$\begin{aligned} \delta\phi \equiv \delta(\Delta\phi(t)) &= \left(\dot{\phi}_{\text{PN}'}(t+t_c-t_m) - \dot{\phi}_{\text{NR}}(t_m)\right) \delta t_m \\ &= \left(\dot{\phi}_{\text{PN}}(t) - \omega_m\right) \delta t_m. \end{aligned} \quad (59)$$

Our simulations (and therefore the comparisons to post-Newtonian theory) start at $m\omega \approx 0.033$, therefore, the maximal error $\delta\phi$ within our comparison at times *before* the matching frequency will be

$$|\delta\phi_{\text{before}}| \leq |0.033 - \omega_m| |\delta t_m| \quad (60)$$

Combining Eqs. (60) and (56), and using $e \approx 6 \times 10^{-5}$, we find that $\delta\phi_{\text{before}} < 0.01$ radians for all four of our matching frequencies $m\omega_m = 0.04, 0.05, 0.063, 0.1$. The maximum error $\delta\phi$ within our comparison at times *after* the matching frequency is

$$|\delta\phi_{\text{after}}| \leq |0.1 - \omega_m| |\delta t_m|, \quad (61)$$

because we end our comparisons to post-Newtonian theory at $m\omega = 0.1$. Eq. (61) evaluates to 0.05 radians for $m\omega_m = 0.04$, and is less than about 0.02 radians for the three higher matching frequencies.

The error in the gravitational wave amplitude caused by an error in t_m can be estimated by Eq. (51). A conservative estimate using $\delta t = 0.9m$ still gives a small error, $\delta A/A \approx 0.004$.

3. Periodic modulation of phase and amplitude

The third effect of orbital eccentricity is a periodic modulation of the gravitational wave phase and amplitude. If we assume that $\bar{\omega}(t)$ varies on much longer time

scales than $1/\Omega_r$ (which is true at large separation) then time integration of Eq. (55) yields

$$\phi(t) = \bar{\phi}(t) + 2e \frac{\bar{\omega}}{\Omega_r} \sin(\Omega_r t). \quad (62)$$

Because $\Omega_r \approx \Omega \approx \bar{\omega}/2$, we therefore find that the gravitational wave phase consists of the sum of the desired “circular” phase, $\bar{\phi}(t)$, plus a oscillatory component with amplitude $4e \approx 2 \times 10^{-4}$. This oscillatory component, however, is much smaller than other uncertainties of the comparison, for instance the uncertainty in determination of t_m .

Residual eccentricity will also cause a modulation of the gravitational wave amplitude in a manner similar to that of the phase. This is because eccentricity explicitly enters the post-Newtonian amplitude formula at 0PN order [131]. This term is proportional to e , and since $e \lesssim 6 \times 10^{-5}$ its contribution to the amplitude error is small compared to the effect due to uncertainty in t_m .

While oscillations in phase and amplitude due to eccentricity are tiny and dwarfed by other uncertainties in the PN-NR comparison, their characteristic oscillatory behavior makes them nevertheless visible on some of the graphs we present below, for instance, both panels of Fig. 18.

F. Spin

We now turn our attention to effects of the small residual spins of the black holes. References [132, 133] compute spin-orbit coupling up to 2.5 post-Newtonian order, and find that the orbital phase, Eq. (40), acquires the following spin contributions

$$\begin{aligned} \Phi_S(x) &= -\frac{1}{32\nu} \sum_{i=1,2} \chi_i \left\{ \left(\frac{565}{24} \frac{m_i^2}{m^2} + \frac{125\nu}{8} \right) x^{-1} \right. \\ &\quad - \left[\left(\frac{681145}{4032} + \frac{965\nu}{28} \right) \frac{m_i^2}{m^2} \right. \\ &\quad \left. \left. + \frac{37265\nu}{448} + \frac{1735\nu^2}{56} \right] \ln x \right\}, \end{aligned} \quad (63)$$

where $\chi_i = \mathbf{S}_i \cdot \hat{\mathbf{L}}/m_i^2$ is the projection of the dimensionless spin of the i -th hole onto the orbital angular momentum. For equal-mass binaries with spins $\chi_1 = \chi_2 \equiv \chi$, this reduces to

$$\Phi_S(x) = -\chi \left(\frac{235}{96} x^{-1} - \frac{270625}{16128} \ln x \right). \quad (64)$$

Our comparisons to post-Newtonian theory are performed over the orbital frequency range of $0.0167 \leq m\Omega \leq 0.05$, corresponding to $0.065 \leq x \leq 0.136$. The gravitational wave phase is approximately twice the orbital phase, so that the spin-orbit coupling contributes

$$\delta\phi_S = 2[\Phi_S(0.065) - \Phi_S(0.136)] \approx -64 \chi \quad (65)$$




FIG. 15: Comparison of numerical simulation with **TaylorT1 3.5/2.5** waveforms. Left: Difference in gravitational wave phase. Right: Relative amplitude difference. Plotted are comparisons for four values of ω_m . The filled diamond on each curve shows the point at which $\dot{\phi} = \omega_m$, and the open circle denotes the point where the post-Newtonian frequency reaches ISCO for a Schwarzschild black hole. The insets show enlargements for small differences and early times. Also shown is the difference between the numerical and restricted (i.e. 3.5PN phase, 0PN amplitude) Taylor T1 for $m\omega_m = 0.1$.

to the gravitational wave phase. In Sec. II B we estimated $|\mathbf{S}|/M_{\text{irr}}^2 < 5 \times 10^{-4}$, where M_{irr} is the irreducible mass of either black hole. Because $\chi \leq |\mathbf{S}|/M_{\text{irr}}^2 \approx 5 \times 10^{-4}$, the residual black hole spins contribute less than 0.03 radians to the overall gravitational wave phase.

We now turn to errors in the amplitude comparison caused by residual spin. From Eq. (64) we can compute the error in orbital frequency as

$$\begin{aligned} \delta\Omega = \dot{\Phi}_s &= \chi \frac{\dot{x}}{x} \left(\frac{235}{96} x^{-1} + \frac{270625}{16128} \right) \\ &= \chi x^4 \frac{16}{5m} \left(\frac{235}{96} x^{-1} + \frac{270625}{16128} \right), \end{aligned} \quad (66)$$

where we have used Eq. (45). Because the amplitude of Ψ_4 scales like $\Omega^{8/3}$, we arrive at

$$\frac{\delta A}{A} = \frac{8}{3} \frac{\delta\Omega}{\Omega} = \chi x^{5/2} \frac{128}{15} \left(\frac{235}{96} x^{-1} + \frac{270625}{16128} \right), \quad (67)$$

which for $m\omega_m = 0.1$ (i.e. $x = 0.136$) gives $\delta A/A = 2.0\chi \sim 1.0 \times 10^{-3}$.

Spin-orbit terms also contribute directly to the amplitude [134, 135]. The leading order contribution (for an equal-mass binary with equal spins) contributes a term $\delta A/A \sim (4/3)\chi x^{3/2}$, which is the same order of magnitude as the previous error, 10^{-3} .

VI. RESULTS

A. Comparison with individual post-Newtonian approximants

We compare our simulations with three different post-Newtonian approximants: the TaylorT1, TaylorT3, and TaylorT4 waveforms. These three waveforms agree with each other up to their respective post-Newtonian expansion orders, but they differ in the way that the uncontrolled higher order terms enter. We start with the comparison to TaylorT1.

1. TaylorT1 (3.5PN phase, 2.5PN amplitude)

Figure 15 compares the numerical simulation to TaylorT1 3.5/2.5 waveforms (i.e. expansion order 3.5PN in phase and 2.5PN in amplitude, the highest expansion orders currently available for generic direction, cf. III B). The left panel shows the phase difference, where we find differences of more than a radian for all four matching frequencies we consider: $\omega_m = 0.04, 0.05, 0.063,$ and 0.01 .

For our largest matching frequency, $m\omega_m = 0.1$, the phase differences are small toward the end of the run by construction. Nevertheless, a phase difference of more than 0.5 radians builds up in the ~ 1.5 cycles after the matching point before the TaylorT1 template generation fails. Recall that $m\omega_m = 0.1$ occurs about 2.2 gravitational wave cycles before our simulations fail, which is still about 1.5 cycles before merger. However, the largest




FIG. 16: Numerical and **TaylorT1 3.5/2.5** waveforms. The PN waveform is matched to the numerical one at $m\omega_m = 0.04$, indicated by the small circle. The lower panel shows a detailed view of the last 10 gravitational wave cycles.

phase disagreement for $m\omega_m = 0.1$ builds up at early times, reaching 1.5 radians at the beginning of our simulation, about 28 cycles before the matching (~ 30 cycles before the end of the simulation), and still showing no sign of flattening even at the start of our simulation.

To achieve phase coherence with the early inspiral waveform, it is therefore necessary to match earlier than $m\omega_m = 0.1$. The left panel of Fig. 15 clearly shows that phase differences at earlier times become smaller when the matching point itself is moved to earlier time. For instance, $m\omega_m = 0.063$ (about eight gravitational wave cycles before the end of our simulation), results in phase differences less than 0.5 radians during the 22 earlier cycles of our evolution. However, the phase difference $\phi_{\text{PN}} - \phi_{\text{NR}}$ does not level off at early times within the length of our simulation, so it seems quite possible that the phase difference may grow to a full radian or more if the numerical simulations could cover many more cycles. We thus estimate that for TaylorT1, to achieve 1-radian phase coherence with the early inspiral may require matching more than 10 cycles before merger. To achieve more stringent error bounds in phase coherence will require matching even earlier: for instance it appears one needs to use $m\omega_m = 0.04$ (about 20 cycles before the end of our simulation) for a phase error of less than $\lesssim 0.1$ radians.

While matching at small ω_m yields good phase coherence early in the run, it produces much larger phase differences late in the run. For example, matching at $m\omega_m = 0.04$ results in a phase difference of almost 2 radians at frequency $m\omega = 0.1$. This rather dramatic disagreement is illustrated in Fig. 16, which plots both

the numerical and the TaylorT1 waveform, matched at $m\omega_m = 0.04$.

The left panel of Fig. 15 also includes a comparison to the so-called restricted TaylorT1 template, where only the leading order amplitude terms are used (*i.e.* 0PN in amplitude). The reason that higher-order amplitude terms affect the phase differences at all is because we are plotting gravitational-wave phase, not orbital phase. However, we see that the effect of these higher-order amplitude terms on the phase difference is small.

We now turn our attention to comparing the amplitudes of the post-Newtonian and numerical waveforms. The right panel of Fig. 15 shows relative amplitude differences between TaylorT1 3.5/2.5 and the numerical waveforms. At early times, the amplitudes agree to within 2 or 3 per cent, the agreement being somewhat better when the matching is performed at early times. At late times, the amplitudes disagree dramatically; a large fraction of this disagreement lies probably in the fact the post-Newtonian point of merger (*i.e.* the point at which the amplitude diverges) occurs at a different time than the numerical point of merger. We also plot the amplitude of the restricted TaylorT1 template. The disagreement between restricted TaylorT1 and the numerical result is much larger, about 5 per cent.

Our comparison with the restricted TaylorT1 waveform (*i.e.* 3.5/0.0) does not agree with the similar comparison of Hannam *et al.* [39]. Ref. [39] finds much smaller phase differences when comparing against waveforms computed with the TaylorT1 implementation in the LIGO Scientific Collaboration Algorithm Library (LAL) [136]. However, at the time the comparison in Ref. [39] was performed the LAL TaylorT1 routine contained several errors which affect the phase of the waveform.⁷ While Ref. [39] states they fixed errors in the post-Newtonian phasing constants due to the errata [119, 120], there were additional errors that may not have been fixed in their code causing discrepancy between their results and ours. All our PN waveforms have been computed using two independent implementations; we have checked that these agree with each other for all post-Newtonian approximants and PN orders we use here, and that our TaylorT1 waveforms agree with the implementation of TaylorT1 in the LAL library, after all LAL code fixes have been implemented.

2. TaylorT3 (3.5PN phase, 2.5PN amplitude)

Figure 17 is the same as Fig.15 except it compares numerical simulations to the TaylorT3 family of wave-

⁷ We have notified the LAL authors of these errors and fixed the code in the LAL software repository on July 7, 2007. We have also confirmed that this subroutine has not been used to generate templates for LSC gravitational wave searches.




FIG. 17: Comparison of numerical simulation with **TaylorT3 3.5/2.5** waveforms. Left: Difference in gravitational wave phase. Right: Relative amplitude difference. Plotted are comparisons for three values ω_m . The filled diamond on each curve shows the point at which $\phi = \omega_m$. The lines end when the frequency of the TaylorT3 waveform reaches its maximum, which happens before $m\omega = 0.1$, so that the matching frequency $m\omega_m = 0.1$ is absent. The left plot also contains TaylorT3 3.0/3.0, matched at $m\omega_m = 0.1$. The insets show enlargements for small differences.

forms. Two differences between TaylorT1 and TaylorT3 are readily apparent from comparing these two figures. The first is that we do not match TaylorT3 3.5/2.5 waveforms at $m\omega_m = 0.1$. This is because the frequency of TaylorT3 3.5/2.5 waveforms reaches a maximum shortly before the formal coalescence time of the post-Newtonian template, and then *decreases*. The maximal frequency is less than 0.1, so that matching at $m\omega_m = 0.1$ is not possible. For this reason, we have also shown in Fig. 17 a comparison with a TaylorT3 3.0/3.0 waveform matched at $m\omega_m = 0.1$. The other major difference between the TaylorT3 3.5/2.5 and TaylorT1 3.5/2.5 comparison is that the phase difference, $\phi_{\text{PN}} - \phi_{\text{NR}}$, has a different sign. While TaylorT1 3.5/2.5 spirals in *more rapidly* than the numerical simulation, TaylorT3 3.5/2.5 *lags behind*. Interestingly, the phase differences from the numerical simulation for both TaylorT1 3.5/2.5 and TaylorT3 3.5/2.5 are of about equal magnitude (but opposite sign). The TaylorT3 3.0/3.0 comparison matched at $m\omega_m = 0.1$ has smaller phase differences than does the TaylorT3 3.5/2.5 comparison, but the slope of the 3.0/3.0 curve in Fig. 17 is nonzero at early times, so it appears that Taylor T3 3.0/3.0 will accumulate significant phase differences at even earlier times, times prior to the start of our simulation. In Fig. 21 it can be seen that matching TaylorT3 3.0/3.0 at $m\omega_m = 0.04$ leads to a good match early, but leads to a phase difference of 0.6 radians by $m\omega = 0.1$.

3. TaylorT4 (3.5PN phase, 2.5PN amplitude)

Figure 18 is the same as Figs. 15 and 17 except it compares numerical simulations to the TaylorT4 PN wave-

forms. The agreement between TaylorT4 waveforms and the numerical results is astonishingly good, far better than the agreement between NR and either TaylorT1 or TaylorT3. The gravitational wave phase difference lies within our error bounds for the entire comparison region $m\omega \leq 0.1$, agreeing to 0.05 radians or better over 29 of 30 gravitational wave cycles. Ref. [37] found agreement between TaylorT4 and their numerical simulation to the level of their numerical accuracy (~ 2 radians), agreeing to roughly 0.5 radians in the gravitational frequency range of $0.054 \leq m\omega \leq 0.1$. Ref. [33] found that NR agrees better with TaylorT4 than with TaylorT1, but the larger numerical errors of these studies did not allow them to see the surprising degree to which NR and TaylorT4 agree. The gravitational wave amplitude of TaylorT4 agrees with the NR waveform to about 1–2 percent at early times, and 8 percent at late times. In Fig. 19 we plot the NR and TaylorT4 waveforms; the two waveforms are visually indistinguishable on the plot, except for small amplitude differences in the final cycles.

On the left panel of Fig. 18 we also show phase comparisons using PN waveforms computed to 3.5PN order in phase but to 0PN and 3.0PN orders in amplitude, for the case $m\omega_m = 0.1$. The PN order of the amplitude expansion affects the phase comparison because we are plotting differences in gravitational-wave phase and not orbital phase. The differences between using 0PN, 2.5PN, and 3.0PN amplitude expansions are evident on the scale of the graph, but because these differences are smaller than our estimated uncertainties (see Table III), we cannot reliably conclude which of these most closely agrees with the true waveform.

Figure 19 shows the waveforms for TaylorT4 3.5/3.0




FIG. 18: Comparison of numerical simulation with **TaylorT4 3.5/2.5** waveforms. Left: Difference in gravitational wave phase. Right: Relative amplitude difference. Plotted are comparisons for four values of ω_m . The filled diamond on each curve shows the point at which $\phi = \omega_m$. The left plot also includes two phase comparisons with expansions of different PN order in amplitude, as labeled.




FIG. 19: Numerical and **TaylorT4 3.5/3.0** waveforms. The PN waveform is matched to the numerical one at $m\omega_m = 0.04$, indicated by the small circle. The lower panel shows a detailed view of the end of the waveform.

and for the numerical simulation. The curves agree to better than the line thickness everywhere, except in the last two cycles.

Figure 20 presents amplitude differences between NR and TaylorT4 as the post-Newtonian order of the amplitude expansion is varied, but the phase expansion remains at 3.5PN. The 2.5PN amplitude curve was already included in the right panel of Fig. 18. We see clearly




FIG. 20: **TaylorT4** amplitude comparison for different PN orders. Shown is the relative difference in gravitational wave amplitude between TaylorT4 and numerical Y_{22} waveforms as a function of time. Matching is performed at $m\omega_m = 0.04$ (indicated by the left vertical grey line, whereas the right vertical grey line indicates $m\omega = 0.1$). All curves use 3.5PN order in phase but different PN orders (as labeled) in the amplitude expansion.

that higher order amplitude corrections generally result in smaller differences. The 3PN amplitude correction to the $l = 2, m = 2$ term recently derived by Kidder [60] improves agreement dramatically over the widely known




FIG. 21: Phase comparison for different PN approximants at different PN orders, matched at $m\omega_m = 0.04$. Shown is the difference in gravitational wave phase between each post-Newtonian approximant and the numerical Y_{22} waveforms as a function of time. The two vertical grey lines indicate when the numerical simulation reaches $m\omega = 0.063$ and 0.1 , respectively; the labels along the top horizontal axes give the number of GW-cycles before $m\omega = 0.1$.

2.5PN amplitude formulae. Unfortunately, the 3PN amplitude correction to the entire waveform, including all Y_{lm} modes, is not known.⁸

B. Comparing different post-Newtonian approximants

The previous section presented detailed comparisons of our numerical waveforms with three different post-Newtonian approximants. We now turn our attention to some comparisons between these approximants. In this section we also explore further how the post-Newtonian order influences agreement between numerical and post-Newtonian waveforms.

Figure 21 presents phase differences as a function of time for all three PN approximants we consider here and for different PN orders. The post-Newtonian and numerical waveforms are matched at $m\omega_m = 0.04$, about 9 cycles after the beginning of the numerical waveform, and about 21 cycles before its end. We find that some PN approximants at some particular orders agree exceed-

⁸ To get the complete waveform to 3PN order, only the (2, 2) mode must be known to 3PN order; other modes must be known to smaller PN orders. For an equal mass, non-spinning binary, all modes except the (3, 2) mode are currently known to sufficient order to get a complete 3PN waveform [60].




FIG. 22: Phase differences between numerical and post-Newtonian waveforms at two selected times close to merger. Waveforms are matched at $m\omega_m = 0.04$, and phase differences are computed at the time when the numerical simulation reaches $m\omega = 0.063$ (left panel) and $m\omega = 0.1$ (right panel). Differences are plotted versus PN order (equal order in phase and amplitude, except the '3.5 PN' points are 3.5/3.0). The 1PN data points are off scale, clustering at -5.5 radians (left panel) and -15 radians (right panel). The thin black bands indicate upper bounds on the uncertainty of the comparison as discussed in Sec. V A.

ingly well with the numerical results. The best match is easily TaylorT4 at 3.5PN order, and the next best match is TaylorT4 at 2.0PN order. Some approximants behave significantly worse, such as the TaylorT1 and TaylorT4 waveforms at 2.5PN order. The 2.5PN and 3PN TaylorT3 waveforms agree very well with the numerical waveform at early times, but at late times they accumulate a large phase difference; the 2.5PN TaylorT3 waveform ends even before the numerical waveform reaches $m\omega = 0.1$ (the rightmost vertical grey line in Fig. 21).

We also find that all three PN approximants, when computed to 3PN order or higher, match the numerical waveform (and each other) quite closely at early times, when all PN approximants are expected to be accurate. However, at late times, $t - r^* > 2500m$, the three PN approximants begin to diverge, indicating that PN is beginning to break down.

Figure 22 presents similar results in a different format. We compute the phase differences between the numerical waveform and the various post-Newtonian approximants at the times when the numerical waveform reaches gravitational wave frequencies $m\omega = 0.063$ and $m\omega = 0.1$ (the times corresponding to these frequencies are also indicated by grey lines in Fig. 21). We then plot these phase differences as a function of the post-Newtonian order (using equal order in phase and amplitude, except for 3.5PN order, where we use 3.0PN in amplitude). The 2.5PN and 3.5PN TaylorT3 and the 2PN TaylorT1 waveforms end before $t_{0.1}$, so these data points cannot be included in the right panel of Fig. 22.

The general trend seen in Fig. 22 is that the phase dif-

ference decreases with increasing PN order. However, this convergence is not monotonic, and the scatter in Fig. 22 can be larger than the phase differences themselves. For example, the 0PN waveforms are about as good as the 2.5PN waveforms for TaylorT1 and TaylorT4, and the 2.5PN TaylorT4 waveform agrees with the numerical results much better than do either the 2PN or 3PN TaylorT4 waveforms. Considering Fig. 22, it seems difficult to make statements about the convergence with PN order for any particular PN approximant, or statements about which PN orders are generally “good”. Given that at fixed PN order the different approximants differ merely by the treatment of uncontrolled higher-order terms, the scatter in Fig. 22 in some sense represents the truncation error at each PN order. While some PN approximants at certain orders may show better agreement with the numerical simulation, we are not aware of any means to predict this besides direct comparisons to numerical simulations (as is done here). In particular, Fig. 22 suggests that the remarkable agreement between our numerical results and the 3.5PN TaylorT4 approximant may be simply due to luck; clearly, more PN-NR comparisons are needed, with different mass ratios and spins, to see if this is the case.

VII. CONCLUSIONS

We have described numerical simulations of an equal mass, non-spinning binary black hole spacetime covering 15 orbits of inspiral just prior to the merger of the two black holes. Using a multi-domain pseudospectral method we are able to extract the gravitational wave content measured by a distant observer with a phase accuracy of better than 0.02 radians over the roughly 30 cycles of gravitational radiation observed. We demonstrate that in order to achieve this accuracy it is necessary to accurately extrapolate the waveform from data obtained at extraction surfaces sufficiently far from the center of mass of the system. When comparing to zero-spin, zero-eccentricity PN formulae, our phase uncertainty increases to 0.05 radians because the numerical simulation has a small but nonzero orbital eccentricity and small but nonzero spins on the holes.

Judging from the case in which we match at $m\omega_m = 0.04$, our numerical simulations are consistent (within our estimated phase uncertainty) with all PN approximants (at the highest PN order) from the beginning of our inspiral until about 15 gravitational wave cycles prior to the merger of the binary. After this point, the various PN approximants begin to diverge, suggesting that the approximation is beginning to break down. Since there are many different PN approximants (including Padé [52] and effective-one-body [34, 50, 56, 57] which were not discussed in this paper) it may be possible to find a clever way to push the PN expansion beyond its breaking point.

Indeed, we find that one approximant, TaylorT4 at 3.5PN in phase, works astonishingly well, agreeing with

our numerical waveforms for almost the entire 30-cycle length of our runs. Given the wide scatter plot of predictions by various PN approximants, it is likely that TaylorT4 3.5/2.5 simply got lucky for the equal mass non-spinning black hole binary. In fact, the assumption of adiabaticity (*i.e.*, circular orbits) is known to lead to much larger phase differences relative to a non-adiabatic inspiral (see Fig. 4 of [54] and [137]) than the phase differences between NR and TaylorT4 we find in Fig. 18. Thus it seems that the uncontrolled higher order terms of TaylorT4 3.5/2.5 balance the error introduced by the adiabaticity assumption to a remarkable degree. It remains to be determined whether similar cancellations occur when the black hole masses are unequal or when the holes have nonzero spin.

Regardless of the robustness of TaylorT4, it seems evident that numerical simulations are needed in order to know which, if any, PN approximant yields the correct waveform after the various approximants begin to diverge. For there is no *a priori* reason why TaylorT4 should be a better choice than TaylorT1 as they differ only in whether the ratio of gravitational wave flux to the derivative of the orbital energy with respect to frequency is left as a ratio of post-Newtonian expansions or re-expanded as a single post-Newtonian expansion.

The surprising accuracy of TaylorT4 in the gravitational frequency range from $m\omega = 0.035$ through $m\omega = 0.15$, for the equal mass, non-spinning inspiral of two black holes, in principle could form a basis for evaluating the errors of numerical simulations. Instead of worrying about errors due to different formulations, initial data, boundary conditions, extraction methods, etc., perhaps a long inspiral simulation could be compared with TaylorT4 3.5/3.0 in order to get a direct estimate of the phase error.

We find that the 3PN contributions to the amplitude of the (2, 2) modes improve their accuracy with respect to the numerical waveforms. This suggests that for accurate parameter estimation, it may be desirable to compute the full 3PN amplitude for the polarization waveforms.

Much work still needs to be done to improve the comparison between NR and PN. Our primary goal is to push our simulations through merger and ringdown so that we may compare various resummed PN approximants and the effective-one-body approximants during the last cycle of inspiral and merger, as well as test TaylorT4 3.5/3.0 closer to merger. We also intend to do long inspirals with arbitrary masses and spins in order to test the robustness of PN over a range of these parameters.

Furthermore we wish to improve our initial data. There is a large amount of ‘junk radiation’ present in the initial data that limits how early we can match PN and NR waveforms. Reduction of this junk radiation [138] would improve the accuracy of our simulations as well.

Finally, we have done just a simple comparison between NR and PN, without including any treatment of effects that are important for real gravitational wave detectors such as limited bandwidth and detector noise. In

order to more directly address the suitability of PN formulae for analyzing data from gravitational wave detectors, it will be necessary to fold in the properties of the detector, to consider specific values for the total mass of the binary, and to fit for the mass from the waveforms rather than assuming that the PN and NR waveforms correspond to the same mass. We leave this for future work.

Acknowledgments

It is a pleasure to acknowledge useful discussions with Stuart Anderson, Alessandra Buonanno, Mark Hannam, Ian Hinder, Luis Lehner, Lee Lindblom, Geoffrey Lovelace, Robert Owen, Oliver Rinne and Kip Thorne. In particular, we would like to thank Rob Owen for estimating the BH spin, Geoffrey Lovelace for his help constructing initial data, Oliver Rinne for providing im-

proved boundary conditions, and Lee Lindblom for his guidance and input throughout this project. This work was supported in part by grants from the Sherman Fairchild Foundation to Caltech and Cornell, and from the Brinson Foundation to Caltech; by NSF grants PHY-0601459, PHY-0652995, DMS-0553302 and NASA grants NAG5-12834, NNG05GG52G at Caltech; by NSF grants PHY-0652952, DMS-0553677, PHY-0652929, and NASA grant NNG05GG51G at Cornell; and by the Z. Smith Reynolds Foundation and NSF grant PHY-0555617 at Wake Forest. We thank NASA/JPL for providing computing facilities that contributed to this work. Some of the simulations discussed here were produced with LIGO Laboratory computing facilities. LIGO was constructed by the California Institute of Technology and Massachusetts Institute of Technology with funding from the National Science Foundation and operates under cooperative agreement PHY-0107417. This paper has been assigned LIGO document number LIGO-P070101-00-Z.

-
- [1] F. Pretorius, *Phys. Rev. Lett.* **95**, 121101 (2005), gr-qc/0507014.
 - [2] F. Pretorius, *Class. Quant. Grav.* **23**, S529 (2006).
 - [3] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, and J. van Meter, *Phys. Rev. Lett.* **96**, 111102 (2006).
 - [4] M. Campanelli, C. O. Lousto, P. Marronetti, and Y. Zlochower, *Phys. Rev. Lett.* **96**, 111101 (2006).
 - [5] B. Brüggmann, J. A. Gonzalez, M. Hannam, S. Husa, U. Sperhake, and W. Tichy, gr-qc/0610128 (2006).
 - [6] P. Diener, F. Herrmann, D. Pollney, E. Schnetter, E. Seidel, R. Takahashi, J. Thornburg, and J. Ventrella, *Phys. Rev. Lett.* **96**, 121101 (2006), gr-qc/0512108.
 - [7] M. Campanelli, C. O. Lousto, and Y. Zlochower, *Phys. Rev. D* **73**, 061501(R) (2006), gr-qc/0601091.
 - [8] P. Marronetti, W. Tichy, B. Brüggmann, J. Gonzalez, M. Hannam, S. Husa, and U. Sperhake, *Class. Quantum Grav.* **24**, S43 (2007), gr-qc/0701123.
 - [9] B. Szilágyi, D. Pollney, L. Rezzolla, J. Thornburg, and J. Winicour, *Class. Quantum Grav.* **24**, S275 (2007), gr-qc/0612150.
 - [10] M. A. Scheel, H. P. Pfeiffer, L. Lindblom, L. E. Kidder, O. Rinne, and S. A. Teukolsky, *Phys. Rev. D* **74**, 104006 (2006), gr-qc/0607056.
 - [11] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, and J. van Meter, *Phys. Rev. D* **73**, 104002 (2006).
 - [12] J. G. Baker, M. Campanelli, F. Pretorius, and Y. Zlochower, *Class. Quantum Grav.* **24**, S25 (2007), gr-qc/0701016.
 - [13] J. A. Gonzalez, U. Sperhake, B. Brüggmann, M. Hannam, and S. Husa, *Phys. Rev. Lett.* **98**, 091101 (2007), gr-qc/0702052.
 - [14] F. Herrmann, I. Hinder, D. Shoemaker, P. Laguna, and R. A. Matzner, *Astrophys. J.* **661**, 430 (2007), gr-qc/0701143.
 - [15] M. Koppitz, D. Pollney, C. Reisswig, L. Rezzolla, J. Thornburg, P. Diener, and E. Schnetter, *Phys. Rev. Lett.* **99**, 041102 (2007), gr-qc/0701163.
 - [16] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, J. R. van Meter, and M. C. Miller, *Astrophys. J.* **653**, L93 (2006), astro-ph/0603204.
 - [17] M. Campanelli, C. O. Lousto, Y. Zlochower, and D. Merritt, *Phys. Rev. Lett.* **98**, 231102 (2007), gr-qc/0702133.
 - [18] J. A. Gonzalez, M. D. Hannam, U. Sperhake, B. Brüggmann, and S. Husa, *Phys. Rev. Lett.* **98**, 231101 (2007), gr-qc/0702052.
 - [19] C. F. Sopuerta, N. Yunes, and P. Laguna, *Astrophys. J.* **656**, L9 (2007), astro-ph/0611110.
 - [20] D.-I. Choi, B. J. Kelly, W. D. Boggs, J. G. Baker, J. Centrella, and J. van Meter (2007), gr-qc/0702016.
 - [21] M. Campanelli, C. O. Lousto, Y. Zlochower, and D. Merritt, *Astrophys. J.* **659**, L5 (2007).
 - [22] B. Brüggmann, J. Gonzalez, M. Hannam, S. Husa, and U. Sperhake (2007), arXiv:0707.0135.
 - [23] J. G. Baker, W. D. Boggs, J. Centrella, B. J. Kelly, S. T. McWilliams, M. C. Miller, and J. R. van Meter (2007), astro-ph/0702390.
 - [24] F. Herrmann, I. Hinder, D. M. Shoemaker, P. Laguna, and R. A. Matzner (2007), arXiv:0706.2541v1 [gr-qc].
 - [25] F. Herrmann, I. Hinderer, D. Shoemaker, and P. Laguna, *Class. Quantum Grav.* **24**, S33 (2007), gr-qc/0601026.
 - [26] J. D. Schnittman, A. Buonanno, J. R. van Meter, J. G. Baker, W. D. Boggs, J. Centrella, B. J. Kelly, and S. T. McWilliams (2007), arXiv:0707.0301v1 [gr-qc].
 - [27] M. Campanelli, C. O. Lousto, Y. Zlochower, and D. Merritt, *Phys. Rev. D* **75**, 064030 (2007), gr-qc/0612076.
 - [28] M. Campanelli, C. O. Lousto, and Y. Zlochower, *Phys. Rev. D* **74**, 084023 (2006), gr-qc/0608275.
 - [29] M. Campanelli, C. O. Lousto, and Y. Zlochower, *Phys. Rev. D* **74**, 041501 (2006), gr-qc/0604012.
 - [30] P. Ajith, S. Babak, Y. Chen, M. Hewitson, B. Krishnan, J. T. Whelan, B. Brüggmann, P. Diener, J. Gonzalez, M. Hannam, et al. (2007), arXiv:0704.3764.
 - [31] E. Berti, V. Cardoso, J. A. Gonzalez, U. Sperhake, M. Hannam, S. Husa, and B. Brüggmann (2007), gr-qc/0703053.

- [32] A. Buonanno, G. B. Cook, and F. Pretorius, *Phys. Rev. D* **75**, 124018 (2007), gr-qc/0610122.
- [33] Y. Pan, A. Buonanno, J. G. Baker, J. Centrella, B. J. Kelly, S. T. McWilliams, F. Pretorius, and J. R. van Meter, arXiv:0704.1964v1 [gr-qc] (2007).
- [34] A. Buonanno, Y. Pan, J. G. Baker, J. Centrella, B. J. Kelly, S. T. McWilliams, and J. R. van Meter (2007), arXiv:0706.3732v2 [gr-qc].
- [35] T. Baumgarte, P. Brady, J. D. Creighton, L. Lehner, F. Pretorius, and R. DeVoe (2006), gr-qc/0612100.
- [36] H. P. Pfeiffer, D. A. Brown, L. E. Kidder, L. Lindblom, G. Lovelace, and M. A. Scheel, *Class. Quantum Grav.* p. S59 (2007).
- [37] J. G. Baker, J. R. van Meter, S. T. McWilliams, J. Centrella, and B. J. Kelly, gr-qc/0612024 (2006).
- [38] J. G. Baker, S. T. McWilliams, J. R. van Meter, J. Centrella, D.-I. Choi, B. J. Kelly, and M. Koppitz, *Phys. Rev. D* **75**, 124024 (2007), gr-qc/0612117.
- [39] M. Hannam, S. Husa, U. Sperhake, B. Brügmann, and J. A. Gonzalez (2007), arXiv:0706.1305v1 [gr-qc].
- [40] S. Husa, J. A. Gonzalez, M. Hannam, B. Brügmann, and U. Sperhake (2007), arXiv:0706.0740v1 [gr-qc].
- [41] B. C. Barish and R. Weiss, *Phys. Today* **52 (Oct)**, 44 (1999).
- [42] S. J. Waldman, *Class. Quantum Grav.* **23**, S653 (2006).
- [43] S. Hild, *Class. Quantum Grav.* **23**, S643 (2006).
- [44] F. Acernese et al., *Class. Quantum Grav.* **19**, 1421 (2002).
- [45] F. Acernese, P. Amico, M. Alshourbagy, F. Antonucci, S. Aoudia, S. Avino, D. Babusci, G. Ballardin, F. Barone, L. Barsotti, et al., *Class. Quantum Grav.* **23**, S635 (2006).
- [46] P. Fritschel, in *Proceedings of SPIE Vol. 4856 "Gravitational Wave Detection"*, edited by M. Cruise and P. Saulson (2003), pp. 282–291, arXiv:gr-qc/0308090v1.
- [47] E. E. Flanagan and S. A. Hughes, *Phys. Rev. D* **57**, 4535 (1998), gr-qc/97101039.
- [48] P. C. Peters, *Phys. Rev.* **136**, B1224 (1964).
- [49] T. Damour, B. R. Iyer, and B. S. Sathyaprakash, *Phys. Rev. D* **57**, 885 (1998).
- [50] A. Buonanno and T. Damour, *Phys. Rev. D* **59**, 084006 (1999).
- [51] A. Buonanno, Y.-b. Chen, and M. Vallisneri, *Phys. Rev. D* **67**, 024016 (2003), erratum *ibid D* **74** 029903 (2006), gr-qc/0205122.
- [52] T. Damour, B. R. Iyer, and B. S. Sathyaprakash, *Phys. Rev. D* **63**, 044023 (2001), erratum-*ibid.D72:029902,2005*.
- [53] T. Damour, B. R. Iyer, and B. S. Sathyaprakash, *Phys. Rev. D* **66** (2002), erratum-*ibid.D72:029901,2005*.
- [54] A. Buonanno and T. Damour, *Phys. Rev. D* **62**, 064015 (2000).
- [55] T. Damour, P. Jaranowski, and G. Schäfer, *Phys. Rev. D* **62**, 084011 (2000), arXiv:gr-qc/0005034.
- [56] T. Damour, *Phys. Rev. D* **64**, 124013 (2001).
- [57] T. Damour, B. R. Iyer, P. Jaranowski, and B. S. Sathyaprakash, *Phys. Rev. D* **67**, 064028 (2003).
- [58] A. Buonanno, Y. Chen, and T. Damour, *Phys. Rev. D* **74**, 104005 (2006).
- [59] S. Husa, M. Hannam, J. A. González, U. Sperhake, and B. Brügmann (2007), arXiv:0706.0904.
- [60] L. E. Kidder, in preparation.
- [61] J. W. York, Jr., *Phys. Rev. Lett.* **82**, 1350 (1999).
- [62] H. P. Pfeiffer and J. W. York, *Phys. Rev. D* **67**, 044022 (2003).
- [63] H. P. Pfeiffer, L. E. Kidder, M. A. Scheel, and S. A. Teukolsky, *Comput. Phys. Commun.* **152**, 253 (2003).
- [64] G. B. Cook, *Phys. Rev. D* **65**, 084003 (2002).
- [65] G. B. Cook and H. P. Pfeiffer, *Phys. Rev. D* **70**, 104016 (2004).
- [66] M. Caudill, G. B. Cook, J. D. Grigsby, and H. P. Pfeiffer, *Phys. Rev. D* **74**, 064011 (2006), gr-qc/0605053.
- [67] L. Lindblom, M. A. Scheel, L. E. Kidder, R. Owen, and O. Rinne, *Class. Quantum Grav.* **23**, S447 (2006).
- [68] F. Pretorius, *Class. Quantum Grav.* **22**, 425 (2005).
- [69] H. Friedrich, *Commun. Math. Phys.* **100**, 525 (1985).
- [70] D. Garfinkle, *Phys. Rev. D* **65**, 044029 (2002).
- [71] O. Rinne, *Class. Quantum Grav.* **23**, 6275 (2006), gr-qc/0606053.
- [72] O. Rinne, L. Lindblom, and M. A. Scheel, *Class. Quantum Grav.* **24**, 4053 (2007).
- [73] J. M. Stewart, *Class. Quantum Grav.* **15**, 2865 (1998).
- [74] H. Friedrich and G. Nagy, *Comm. Math. Phys.* **201**, 619 (1999).
- [75] J. M. Bardeen and L. T. Buchman, *Phys. Rev. D* **65**, 064037 (2002).
- [76] B. Szilágyi, B. Schmidt, and J. Winicour, *Phys. Rev. D* **65**, 064015 (2002).
- [77] G. Calabrese, J. Pullin, O. Sarbach, M. Tiglio, and O. Reula, *Commun. Math. Phys.* **240**, 377 (2003).
- [78] B. Szilágyi and J. Winicour, *Phys. Rev. D* **68**, 041501(R) (2003).
- [79] L. E. Kidder, L. Lindblom, M. A. Scheel, L. T. Buchman, and H. P. Pfeiffer, *Phys. Rev. D* **71**, 064020 (2005).
- [80] L. T. Buchman and O. C. A. Sarbach, *Class. Quantum Grav.* **23**, 6709 (2006).
- [81] M. Björhus, *SIAM J. Sci. Comput.* **16**, 542 (1995).
- [82] D. Gottlieb and J. S. Hesthaven, *J. Comp. Appl. Math.* **128**, 83 (2001).
- [83] J. S. Hesthaven, *Appl. Numer. Math.* **33**, 23 (2000).
- [84] J. D. Brown and J. W. York, *Phys. Rev. D* **47**, 1407 (1993).
- [85] A. Ashtekar, C. Beetle, and J. Lewandowsky, *Phys. Rev. D* **64**, 044016 (2001).
- [86] A. Ashtekar and B. Krishnan, *Phys. Rev. D* **68**, 104030 (2003).
- [87] O. Dreyer, B. Krishnan, E. Schnetter, and D. Shoemaker, *Phys. Rev. D* **67**, 024018 (2003).
- [88] G. B. Cook and B. F. Whiting, *Phys. Rev. D* **76**, 041501(R) (2007).
- [89] R. Owen, Ph.D. thesis, California Institute of Technology (2007), <http://resolver.caltech.edu/CaltechETD:etd-05252007-143511>.
- [90] D. R. Fiske, J. G. Baker, J. R. van Meter, D.-I. Choi, and J. M. Centrella, *Phys. Rev. D* **71**, 104036 (2005).
- [91] C. Beetle, M. Bruni, L. M. Burko, and A. Nerozzi, *Phys. Rev. D* **72**, 024013 (2005).
- [92] A. Nerozzi, C. Beetle, M. Bruni, L. M. Burko, and D. Pollney, *Phys. Rev. D* **72**, 024014 (2005).
- [93] L. M. Burko, T. W. Baumgarte, and C. Beetle, *Phys. Rev. D* **73**, 024002 (2006).
- [94] M. Campanelli, B. J. Kelly, and C. O. Lousto, *Phys. Rev. D* **73**, 064005 (2006).
- [95] A. Nerozzi, M. Bruni, L. M. Burko, and V. Re, gr-qc/0607066 (2006).
- [96] E. Pazos, E. N. Dorband, A. Nagar, C. Palenzuela, E. Schnetter, and M. Tiglio (2006), gr-qc/0612149.

- [97] L. Lehner and O. M. Moreschi, ArXiv e-prints **706** (2007), 0706.1319.
- [98] B. Kocsis and A. Loeb, *Distortion of gravitational-wave packets due to their self-gravity* (2007), URL <http://arxiv.org/abs/0704.1149>.
- [99] L. Blanchet, Living Rev. Relativity **9**, 4 (2006).
- [100] P. Jaranowski and G. Schäfer, Phys. Rev. D **57**, 7274 (1998), erratum-ibid.D63:029902,2001.
- [101] P. Jaranowski and G. Schäfer, Phys. Rev. D **60**, 124003 (1999).
- [102] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **62**, 021501 (2000), erratum-ibid.D63:029903,2001.
- [103] T. Damour, P. Jaranowski, and G. Schäfer, Phys. Rev. D **63**, 044021 (2001), erratum-ibid.D66:029901,2002.
- [104] L. Blanchet and G. Faye, Physics Letters A **271**, 58 (2000).
- [105] L. Blanchet and G. Faye, Phys. Rev. D **63**, 062005 (2001).
- [106] T. Damour, P. Jaranowski, and G. Schäfer, Physics Letters B **513**, 147 (2001).
- [107] L. Blanchet, T. Damour, and G. Esposito-Farèse, Phys. Rev. D **69**, 124007 (2004).
- [108] Y. Itoh, T. Futamase, and H. Asada, Phys. Rev. D **63**, 064038 (2001).
- [109] Y. Itoh and T. Futamase, Phys. Rev. D **68**, 121501 (2003).
- [110] Y. Itoh, Phys. Rev. D **69** (2004), 064018.
- [111] L. Blanchet and B. R. Iyer, Classical and Quantum Gravity **20**, 755 (2003).
- [112] V. C. de Andrade, L. Blanchet, and G. Faye, Classical and Quantum Gravity **18**, 753 (2001).
- [113] K. S. Thorne, Rev. Mod. Phys. **52**, 299 (1980).
- [114] L. Blanchet, Classical and Quantum Gravity **15**, 1971 (1998).
- [115] L. Blanchet and T. Damour, Phys. Rev. D **46**, 4304 (1992).
- [116] L. Blanchet, Classical and Quantum Gravity **15**, 89 (1998).
- [117] L. Blanchet, Classical and Quantum Gravity **15**, 113 (1998), erratum-ibid.22:3381,2005.
- [118] C. Cutler, T. A. Apostolatos, L. Bildsten, L. S. Finn, E. E. Flanagan, D. Kennefick, D. M. Markovic, A. Ori, E. Poisson, and G. J. Sussman, Physical Review Letters **70**, 2984 (1993).
- [119] L. Blanchet, B. R. Iyer, and B. Joguet, Phys. Rev. D **65**, 064005 (2002), erratum-ibid.D71:129903,2005.
- [120] L. Blanchet, G. Faye, B. R. Iyer, and B. Joguet, Phys. Rev. D **65** (2002), erratum-ibid.D71:129902,2005.
- [121] L. Blanchet and B. R. Iyer, Phys. Rev. D **71**, 024004 (2005).
- [122] L. Blanchet, T. Damour, G. Esposito-Farèse, and B. R. Iyer, Physical Review Letters **93**, 091101 (2004).
- [123] L. Blanchet, T. Damour, G. Esposito-Farèse, and B. R. Iyer, Phys. Rev. D **71**, 124004 (2005).
- [124] K. Arun, L. Blanchet, B. Iyer, and M. Qusailah, Class. Quantum Grav. **21**, 3771 (2004), erratum-ibid. 22, 3115–3117 (2005).
- [125] L. E. Kidder, L. Blanchet, and B. R. Iyer, arXiv:0706.0726 [gr-qc] (2007).
- [126] A. G. Wiseman, Phys. Rev. D **48**, 4757 (1993).
- [127] L. Blanchet and G. Schäfer, Class. Quant. Grav. **10**, 2699 (1993).
- [128] L. Blanchet, B. R. Iyer, C. M. Will, and A. G. Wiseman, Classical and Quantum Gravity **13**, 575 (1996).
- [129] L. Blanchet, Phys. Rev. D **55**, 714 (1997).
- [130] A. Królak, K. D. Kokkotas, and G. Schäfer, Phys. Rev. D **52**, 2089 (1995).
- [131] H. Wahlquist, General Relativity and Gravitation **19**, 1101 (1987).
- [132] G. Faye, L. Blanchet, and A. Buonanno, Phys. Rev. D **74**, 104033 (2006), erratum **75**, 049903(E) (2007).
- [133] L. Blanchet, A. Buonanno, and G. Faye, Phys. Rev. D **74**, 104034 (2006), erratum **75**, 049903(E) (2007).
- [134] L. E. Kidder, Phys. Rev. D **52**, 821 (1995).
- [135] C. M. Will and A. G. Wiseman, Phys. Rev. D **54**, 4813 (1996).
- [136] LSC Algorithm Library software packages LAL, LALWRAPPER, and LALAPPS, URL <http://www.lsc-group.phys.uwm.edu/lal>.
- [137] M. Miller, Phys. Rev. D **69**, 124013 (2004).
- [138] G. M. Lovelace, Ph.D. thesis, California Institute of Technology (2007), <http://etd.caltech.edu/etd/available/etd-05232007-115433>.